

Succinct Certificates for the Solvability of Binary Quadratic Diophantine Equations

J. C. Lagarias

Department of Mathematics
University of Michigan
Ann Arbor, MI 48109-1043
lagarias@umich.edu

(November 6, 2006)

Abstract

Binary quadratic Diophantine equations are of interest from the viewpoint of computational complexity theory. This class of equations includes as special cases many of the known examples of natural problems apparently occupying intermediate stages in the $P - NP$ hierarchy, i.e., problems not known to be solvable in polynomial time nor to be NP -complete, for example the problem of factoring integers.

Let $L(F)$ denote the length of the binary encoding of the binary quadratic Diophantine equation F given by $ax_1^2 + bx_1x_2 + cx_2^2 + dx_1 + ex_2 + f = 0$. Suppose F is such an equation having a nonnegative integer solution. This paper shows that there is a proof (i.e., “certificate”) that F has such a solution which can be checked in $O(L(F)^5 \log L(F) \log \log L(F))$ bit operations. A corollary of this result is that the set $\Sigma = \{F : F \text{ has a nonnegative integer solution}\}$ is in the complexity class NP . The result that Σ is in NP is interesting because it is known that there are binary quadratic Diophantine equations whose smallest nonnegative integer solution is so large that it requires time exponential in $L(F)$ just to write this solution down in the usual binary representation.

1. Introduction

There has been considerable interest in bounding the computational complexity of various number-theoretic problems. A particular motivation is the Rivest-Shamir-Adleman enciphering scheme [31] whose resistance to cryptanalysis depends on the apparent difficulty of factoring large integers. Many of these number-theoretic problems can be formulated as one of two types of problems involving Diophantine equations.

- (i) Deciding whether a given Diophantine equation has an admissible solution or not.
- (ii) Exhibiting an admissible solution to such an equation when it has one.

Here an *admissible* solution denotes an integer solution which may also be required to satisfy some side conditions characteristic of the particular problem. The side conditions that arise are generally of the following two types:

- (i) *Nonnegativity*. Certain variables are required to be nonnegative.

- (ii) *Congruence.* Certain variables x_i are required to satisfy congruence restrictions $x_i \equiv \alpha_i \pmod{\Gamma}$ where the α_i and Γ are given as input.

In this framework, for example, an integer N is composite if and only if the binary quadratic Diophantine equation

$$(x+2)(y+2) = N \quad (1.1)$$

has a solution in nonnegative integers. The problem of factoring N involves exhibiting nonnegative solutions to a series of equations (1.1), and showing that certain other equations of the form (1.1) are solvable.

There is a close relation between Diophantine equations and the theory of computation. The methods developed by Davis, Putman, Robinson and Matijasevic in their solution to Hilbert's 10th problem established that for any recursively enumerable set A of natural numbers there is a Diophantine equation $P(x_1, \dots, x_n) = 0$ such that

$$x \in A \Leftrightarrow \exists \text{ nonnegative } x_2, \dots, x_n \text{ such that } P(x, x_2, \dots, x_n) = 0$$

(see in particular [12], [13], [26].) Adleman and Manders [1], [2] used these methods to establish a computational complexity theory based on the notion of recognizing sets for which a given Diophantine equation has solutions of size bounded by a given complexity function Φ . More precisely, they considered sets S given

$$x \in S \Leftrightarrow \exists \text{ nonnegative } x_2, \dots, x_n \text{ with } L(x_2), \dots, L(x_n) \leq \Phi(L(x))$$

$$\text{such that } P(x, x_2, \dots, x_n) = 0$$

where $P(x_1, \dots, x_n)$ is a fixed Diophantine equation, $L(x)$ denotes the length of the binary integer x , and $\Phi(t)$ is a complexity measure which is an increasing function of t . They introduced a complexity class D which is a Diophantine analogue of the complexity class NP . It consists of all relations $R \subseteq \omega^m$ specified by a Diophantine equation $P(x_1, \dots, x_{m+n}) = 0$ and a polynomial $q(t)$ as follows:

$$\begin{aligned} \langle x_1, \dots, x_m \rangle \in R &\Leftrightarrow \exists \text{ nonnegative } y_1, \dots, y_n \\ &\text{such that } \max L(y_1) \leq q(L(x_1) + \dots + L(x_m)) \\ &\text{and } P(x_1, \dots, x_m, y_1, \dots, y_n) = 0. \end{aligned} \quad (1.2)$$

It is immediately clear that $D \subseteq NP$. It is an open problem whether or not $D = NP$; this is an important problem in determining the relative computing power of Diophantine equations as compared to that of nondeterministic Turing machines.

The class of *binary quadratic Diophantine equations*

$$ax_1^2 + bx_1x_2 + cx_2^2 + dx_1 + ex_2 + f = 0 \quad (1.3)$$

is of special interest from the viewpoint of both number theory and complexity theory. From the viewpoint of number theory, this class of equations can be used to encode the problem of factorization, of solving quadratic congruences

$$x^2 = f \pmod{e}$$

(which corresponds to $x_1^2 - ex_2 - f = 0$), of solving Pell's equation

$$x_1^2 - dx_2^2 = 1 ,$$

and to encode problems in representation and equivalence of binary quadratic forms. From the viewpoint of complexity theory this class of equations seems to represent the borderline between tractable and intractable computational problems. It includes as special cases most of the known examples of natural problems apparently occupying intermediate stages in the $P - NP$ hierarchy (i.e., problems not known to be in P not to be NP -complete) as well as NP -complete problems. For example, it known that

- (i) $S = \{p | p \text{ is prime}\} \in P$ (Agrawal, Kayal and Saxena [6]).
- (ii) $\{\alpha, \beta, \gamma \in \omega | \exists \text{ nonnegative } x_1, x_2 \text{ such that } ax_1^2 + \beta x_2 - \gamma = 0\}$ is NP -complete. (Manders and Adleman [24]).
- (iii) $\{a, c \in \omega | \exists x_1, x_2 \text{ such that } ax_1x_2 + x_2 = c\} \in NP \setminus NP^c$ provided $NP \neq NP^c$. This problem is γ -complete (Adleman and Manders [3]).
- (iv) $\{a, c \in \omega | \exists x_1, x_2 \text{ such that } x_1^2 - a^2x_2^2 = c\}$ is unfaithfully random complete (Adleman and Manders [4], [5]).

All of the sets (i)–(iv) are in D , and hence are certainly in NP . Note that the existence of NP -complete sets in D does not establish $D = NP$. Indeed, Adleman and Manders [2] exhibit a set in P not known to be in D .

This paper treats the general problem of recognizing those binary quadratic Diophantine equations which have nonnegative solutions, which may also be required to satisfy given congruence side conditions. This problem appears to be fundamentally harder computationally than any of the special subclasses of binary quadratic Diophantine equations considered up to now (i.e. including (i)–(iv) above) as indicated by the following example.

Example. Consider the set of equations

$$x^2 - dy^2 = -1 \tag{1.4}$$

where d is given in its binary representation as input. The equations (1.4) are often called the *non-Pellian* or *anti-Pellian* equations. For the subset $d = 5^{2n+1}$, the input requires no more than $7n$ bits. In Lagarias [21, Appendix A], it is shown that for $d = 5^{2n+1}$ this equation has solutions for $n = 1, 2, 3, \dots$ and that the solution (t_1, u_1) to this equation with minimal binary lengths $L(t_1), L(u_1)$ is given by

$$t_1 + u_1\sqrt{5} = (2 + \sqrt{5})^{5^n} . \tag{1.5}$$

This implies that the length of *any* solution x to (1.4) expressed in binary for these d satisfies

$$L(x) > \frac{1}{3}5^n . \quad \square$$

This example shows that there are some binary quadratic Diophantine equations whose solutions are so large that it requires exponential space (in terms of the length of the coefficients of the equation) to store any such solution in binary. In addition this shows the set

$$\Pi = \{d \in \omega \mid x^2 - dy^2 = -1 \text{ is solvable in integers} \} \quad (1.6)$$

cannot be established to be in the complexity class D by use of the Diophantine equation $x^2 - dy^2 + 1 = 0$ in (1.2). Indeed we cannot hope to show the set (1.6) is in NP by guessing a solution x, y , because it potentially requires exponential time to check that a given solution x, y to (1.4) is a solution. (However, see Theorem 1.3 below.) These results apply to the general quadratic Diophantine equation (1.3), because the problem of recognizing the subclass (1.4) is clearly in P .

The main result of this paper is that there exist short certificates of the solvability of all binary quadratic Diophantine equations which have solutions. By the preceding example, these certificates must sometimes verify that solutions exist without exhibiting these solutions in binary. The certificates actually contain in a compact form enough information to exactly calculate an admissible solution.

In order to state the main result, we need two definitions. Given a binary quadratic Diophantine equation $F(x_1, x_2) = 0$ where

$$F(x_1, x_2) = ax_1^2 + bx_1x_2 + cx_2^2 + dx_1 + ex_2 + f$$

together with a congruence side condition

$$\begin{aligned} x_1 &\equiv \alpha_1 \pmod{\Gamma} \\ x_2 &\equiv \alpha_2 \pmod{\Gamma} \end{aligned}$$

and $0 \leq \alpha_1, \alpha_2 < \Gamma$ we define the *length* $L(F)$ of the input to be

$$L(F) = L(a) + L(b) + L(c) + L(d) + L(e) + L(f) + 3L(\Gamma) . \quad (1.7)$$

The other definition concerns the measurement of the running time of a program. We shall measure running in terms of *elementary operations*, which consists of a Boolean operation on a bit or pair of bits, and an input or shift of a single bit. Our main result is the following.

Theorem 1.1. *Let $F(x_1, x_2) = 0$ be a binary quadratic Diophantine equation, where*

$$F(x_1, x_2) = ax_1^2 + bx_1x_2 + cx_2^2 + dx_1 + ex_2 + f ,$$

which has a nonnegative integer solution (x_1, x_2) satisfying

$$\begin{aligned} x_1 &\equiv \alpha_1 \pmod{\Gamma} \\ x_2 &\equiv \alpha_2 \pmod{\Gamma} . \end{aligned}$$

Then there exists a certificate showing that $F(x_1, x_2) = 0$ has such an admissible solution which requires $O(L(F)^5 \log L(F) \log \log L(F))$ elementary operations to verify.

This result gives certificates imposing two side conditions on the solutions: a congruence condition and a positivity condition. The theorem is formulated imposing a positivity side condition to give a result compatible with the framework of Hilbert's 10-th problem, and with Diophantine complexity theory of Adleman and Manders [3], [4]. An immediate consequence of the form of the certificates produced by Theorem 1.1 is the following result.

Theorem 1.2. *The following sets Σ_i are all in NP.*

- (i) $\Sigma_1 = \{a, b, c, d, e, f, \alpha_1, \alpha_2, \Gamma \in \mathbb{Z} \mid \exists \text{ nonnegative integers } x_1, x_2 \text{ with } x_1 \equiv \alpha_1 \pmod{\Gamma}, x_2 \equiv \alpha_2 \pmod{\Gamma} \text{ and } ax_1^2 + bx_1x_2 + cx_2^2 + dx_1 + ex_2 + f = 0\}$
- (ii) $\Sigma_2 = \{a, b, c, d, e, f \in \mathbb{Z} \mid \exists \text{ nonnegative integers } x_1, x_2 \text{ with } ax_1^2 + bx_1x_2 + cx_2^2 + dx_1 + ex_2 + f = 0\}$
- (iii) $\Sigma_3 = \{a, b, c, d, e, f \in \mathbb{Z} \mid \exists \text{ integers } x_1, x_2 \text{ with } ax_1^2 + bx_1x_2 + cx_2^2 + dx_1 + ex_2 + f = 0\}$.

Since one can tell in polynomial time whether or not a binary quadratic Diophantine equation (1.3) is of the special form

$$ax_1^2 + ex_2 + f = 0 \tag{1.8}$$

and since the set of equations of the form (1.8) which have nonnegative integral solutions is NP-complete [24], we conclude that: *testing membership in the sets Σ_1 and Σ_2 in Theorem 1.2 are both NP-complete.* We are unable to decide whether or not any of the sets Σ_i above are in the Diophantine complexity class D .

The proofs of Theorems 1.1 and 1.3 use the theory of binary quadratic forms as developed by Gauss in *Disquisitiones Arithmeticae* [15]. A treatment of this theory can be found in Buell [8]. The certificates are based on Gauss' operation of *composition* of forms, and crucially use an idea of Shanks [33], which he called the "infrastructure" of quadratic forms. Shanks did not give detailed proofs, but the "infrastructure" method was put on a rigorous footing by Lenstra [23] in 1980, in the framework of quadratic number fields. This paper gives an alternate justification of the infrastructure method in the framework of composition of forms (Lemma 6.2).

In addition to the main theorem, we show that whenever two integer binary quadratic forms are equivalent there exist succinct certificates verifying this equivalence (Theorem 7.1).

We add some remarks on related work. The following result is a direct consequence of [21, Theorem 1.1].

Theorem 1.3. *The set*

$$\Pi = \{d \mid \exists \text{ integers } x, y \text{ with } x^2 - dy^2 = -1\}$$

is in $NP \cap co-NP$.

The problem of characterizing the set Π has been extensively studied in algebraic number theory, see Narkiewicz [28, pp. 124–126], and Redei [30].

In another direction Gurari and Ibarra [16] consider a class of Diophantine equations containing (1.8) as a special case, but not containing (1.3). They show that the subclass of such equations having nonnegative solutions is in D , i.e., when such equations have a nonnegative solution, they have one that is small enough to serve as a certificate. Their NP-completeness result then follows from [24].

The proof of Theorem 1.1 is outlined in Section 2, and the details appear in the following sections.

The results of this paper were presented at the 1979 FOCS conference [19]. A complete paper appeared as the 1981 technical report [22]. The present paper differs from this technical report mainly in the addition and update of references. Our impetus to put this paper on record, after a long delay, is its relevance to Problem 5 of the mathematical problems for the twenty-first century formulated in 2000 by S. Smale [34].

We remark on related developments. The "infrastructure" method of Shanks was by Buchmann and Williams [7] to give succinct certificates for class numbers and approximate representation of regulators of quadratic number fields, under the assumption of the generalized Riemann hypothesis. In 1994 Theil [36] showed that verifying the class number falls in the class $\text{NP} \cap \text{co-NP}$, assuming the generalized Riemann hypothesis. The infrastructure method is well known to be computationally effective in practice, as described in Chapter 5 of Cohen [11], and has been used in computations of class numbers and regulators of quadratic and cubic number fields and function fields.

Acknowledgment. This work was done while the author visited the University of Maryland in 1978-1979, and at AT &T Bell Laboratories in 1979-1981.

2. Outline of the Proof

In this section we describe the main ideas of the proof and establish some notational conventions.

We deal throughout with a system F consisting of a binary quadratic Diophantine equation with side conditions having the form:

$$ax_1^2 + 2bx_1x_2 + cx_2^2 + 2dx_1 + 2ex_2 + f = 0, \quad (2.1)$$

$$x_i \equiv \alpha_i \pmod{\Gamma}, \quad i = 1, 2, \quad (2.2)$$

$$x_i \geq 0, \quad i = 1, 2, \quad (2.3)$$

in order to maintain compatibility with Gauss' formulation of this problem. Any system can be brought to this form by multiplying (1.3) by 2. A solution to (2.1)–(2.3) will be called *admissible*.

Binary quadratic Diophantine equations (2.1) are classified as *definite*, *indefinite* or *degenerate* according as the *determinant*

$$D = b^2 - ac \quad (2.4)$$

is negative, positive and not a square, or a perfect square, respectively. This classification is useful because the sets of solutions to these three types of equations have qualitatively different characteristics. For example, definite and degenerate binary quadratic Diophantine equations always have admissible solutions small enough to serve directly as certificates, if they have any admissible solutions at all (Lemma 3.2). The body of the proof concerns the case of indefinite binary quadratic Diophantine equations.

Gauss [15, Art. 216–221] gave a method to determine whether (2.1) has any integer solutions and if so to give a complete parametric description of all solutions. This method is based on his theory of integral binary quadratic forms, and in particular on determining the equivalence or inequivalence of such forms. Gauss' method easily extends to include the side condition

(2.2), but (2.3) adds new complications. We follow the outline of Gauss' method in reducing the problem to that of recognizing the equivalence of two quadratic forms. In Section 3 we transform the problem to that of studying the equation

$$x_1^2 - Dy_2^2 = g$$

with side conditions. In Section 4 quadratic forms are introduced and the problem is transformed to that of demonstrating that the reduced identity form $\tilde{I}$ of determinant D is equivalent to a reduced form Q_{red} via an equivalence matrix W having certain properties. (See Section 4 for definitions.) The proof are complicated by the need to bound the size of the least admissible solution and to keep track of the nonnegativity condition (2.3) under these transformations.

These reductions have not yet addressed the main difficulty in finding certificates of solvability, which is the possible exponentially large size of the number of binary bits in the least admissible solution. This difficulty is transformed into the possible equally large size of the entries of the equivalence matrix W . In Lemma 4.2 we show that in order to verify admissibility we need only know the entries of W to sufficient accuracy to check a certain sign condition, and that these entries satisfy certain congruence side conditions.

The remainder of the proof is devoted to a detailed study of equivalence matrices showing the equivalence of the reduced identity form $\tilde{I}$ and reduced forms Q_{red} . In Section 5 we describe results of Gauss. Gauss defined a notion of two reduced forms being neighbors. If we form a graph in which the reduced forms are vertices, and edges correspond to two reduced forms being neighbors, then Gauss showed this graph is a union of disjoint cycles, and the cycle including $\tilde{I}$ contains exactly the reduced forms equivalent to $\tilde{I}$, which we call the principal cycle. These results imply that the associated equivalence matrices have a very special form, which is related to the ordinary continued fraction algorithm. However this form alone is insufficient to produce succinct certificates.

The main device of the proof uses another set of relations between these equivalence matrices, which come from Gauss' operation of composition of binary quadratic forms (Lemma 6.1). The idea of using the action of composition of forms on the principal cycle is due to Shanks [33], who called it the "infrastructure". The "infrastructure" asserts that composition is a kind of doubling of distance on the graph of the set of forms. Shanks did not give detailed proofs, but a rigorous justification of the "infrastructure" was given by Lenstra [23], in the language of ideals. In this paper we rigorously justify it in the language of composition of forms, in Lemma 6.2. The action of composition can be combined with Gauss' to find a short sequence of composition formulae that prove the equivalence of any two given forms in the principal cycle (Lemma 6.3). In effect each composition causes a squaring, so that if one multiplied out all the compositions to write down the matrix giving the equivalence, the resulting entries would potentially have exponentially many digits, in terms of the input size. However the correctness of the composition can be checked in each formula separately, avoiding this potential exponential blowup. In Section 7 we further apply these formulae to give succinct certificates for the equivalence of two equivalent indefinite binary quadratic forms. We remark that the equivalence or inequivalence of two definite or degenerate quadratic forms can be decided in polynomial time [20].)

In Section 8 we complete the proof by showing that the formulae of Lemma 6.2 can be used to check that the entries of W satisfy a *given side congruence condition*, and to evaluate W using floating-point computations to enough accuracy to verify a *sign condition on the solutions*. This is of interest if one wishes to recognize positive solutions, as are studied in Hilbert's 10-th

problem. It requires significant extra work to establish these extra side condition properties. In general it is difficult to rigorously prove results that the number of significant digits present after a sequence of floating-point operations, because there is the possibility of losing all significant figures when adding two nearly equal floating-point number of opposite signs. We are able to show this potential cancellation effect cannot occur here, using a priori information about the magnitudes of the quantities being computed at all intermediate steps of the computation.

Appendix A gives bounds on the period lengths (mod M) of solutions to certain second-order linear recurrences. Appendix B gives needed results on floating point computation, concerning bounds on the loss of accuracy in floating-point operations.

2.1. Notations and Conventions.

We use a notion of *size* $\|F\|$ of the system (2.1)–(2.3) given by

$$\|F\| = \text{MAX}(|a|, |b|, |c|, |d|, |e|, |f|, |\Gamma|) . \quad (2.5)$$

The size $\|F\|$ is related to the length $L(F)$ in (??) by

$$\log \|F\| \leq L(F) \leq \log \|F\| + 4 . \quad (2.6)$$

We also need an analogous notion of *size* $\|\mathbf{M}\|$ of a matrix $\mathbf{M} = [m_{ij}]$ given by

$$\|\mathbf{M}\| = \text{MAX}_{i,j} |m_{ij}| . \quad (2.7)$$

If M, N are $m \times k$ and $k \times n$ matrices, respectively, then we have the trivial bound

$$\|\mathbf{MN}\| \leq k \|\mathbf{M}\| \|\mathbf{N}\| . \quad (2.8)$$

For ease in counting operations, we establish the convention that

$$\log x = \begin{cases} \log_2 |x| & |x| \geq 4 \\ 2 & |x| < 4 . \end{cases} \quad (2.9)$$

In addition, when counting operations we will sometimes use the function

$$M(n) = n(\log n)(\log \log n) \quad (2.10)$$

arising from the Schonhage-Strassen bound $O(M(n))$ for the multiplication of two n bit binary integers. The O -symbol has the usual meaning, that $O(f(n))$ means $\leq c|f(n)|$, where c is an absolute, effectively computable positive constant (which may differ at each occurrence of the O -symbol.)

3. Bounding the Size of the Least Admissible Solution

In this section we bound the size of the least admissible solution to definite and degenerate binary quadratic Diophantine equations, and show that this solution itself may serve as a certificate. The difficult case is that of indefinite binary quadratic Diophantine equations. In this case we obtain an exponential upper bound for the size of the minimal admissible solution.

We can immediately simplify (2.1) by an invertible variable change provided $D \neq 0$, $c \neq 0$. Following Mathews ([25], 258–260) we introduce new variables

$$\begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} D & 0 \\ b & c \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} be - cd \\ e \end{bmatrix} \quad (3.1)$$

which transforms (2.1) to the equation

$$y_1^2 - Dy_2^2 = g, \quad (3.2)$$

where

$$g = -c \begin{vmatrix} a & b & c \\ b & c & e \\ d & e & f \end{vmatrix}. \quad (3.3)$$

We note the bound

$$|g| < 6||F||^4. \quad (3.4)$$

Inverting the system (3.1) yields

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} \frac{1}{D} & 0 \\ -\frac{b}{cD} & \frac{1}{c} \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} + \begin{bmatrix} \frac{cd - be}{D} \\ \frac{ae - bd}{D} \end{bmatrix}. \quad (3.5)$$

We have obtained the following.

Lemma 3.1. *Let (y_1, y_2) be an integral solution to (3.2), and let a modulus Γ be given. Then (x_1, x_2) given by (3.5) is a rational solution to (2.1). The congruence class of $(y_1, y_2) \pmod{cD\Gamma}$ determines whether x_1, x_2 is integral, and if so specifies $(x_1, x_2) \pmod{\Gamma}$.*

We now bound the size of solutions to define and degenerate binary quadratic Diophantine equations.

Lemma 3.2. *Suppose that a given binary quadratic Diophantine equation is either definite or degenerate. If it has any admissible solutions at all, then it has an admissible solution (x_1, x_2) with*

$$\text{MAX}(\log x_1, \log x_2) \leq 8 \log ||F|| + 8. \quad (3.6)$$

Proof. There are several cases.

Case 1. $D \neq 0$, $c \neq 0$. Then the change of variables (3.1) takes integral solutions of (2.1) to integral solutions of

$$y_1^2 - Dy_2^2 = g. \quad (3.7)$$

In the definite case $D < 0$, hence all integer solutions to (3.7) have

$$\text{MAX}(|y_1|, |y_2|) \leq \sqrt{|g|}.$$

Using (3.4) and (3.5), we see that all integer solutions to (2.1) must have

$$MAX(|x_1|, |x_2|) \leq ||F||^4 .$$

This implies (3.6). In the degenerate case $D = h^2$ is a perfect square and (3.7) becomes

$$(y_1 + hy_2)(y_1 - hy_2) = g . \quad (3.8)$$

Hence each integer solution (y_1, y_2) gives rise to a factorization $g = g_1 g_2$ where

$$\begin{aligned} y_1 + hy_2 &= g_1 \\ y_1 - hy_2 &= g_2 . \end{aligned} \quad (3.9)$$

Solving (3.9) for y_1, y_2 we obtain

$$MAX(|y_1|, |y_2|) < |g|$$

using (3.5) again, this implies (3.6).

Case 2. $D \neq 0, c = 0, a \neq 0$. Interchange x_1 and x_2 , proceed as in Case 1.

Case 3. $D \neq 0, a = 0, c = 0$. Then (2.1) yields

$$(bx_1 + e)(bx_2 + d) = de - \frac{bf}{2} .$$

A similar argument to (3.8), (3.9) shows

$$MAX(|x_1|, |x_2|) \leq 3||F||^2$$

in this case, implying (3.6).

Case 4. $D = 0$.

$$ax_1^2 + 2bx_1x_2 + cx_2^2 = m(ax_1 + \beta x_2)^2$$

where m, α, β are integers, $m \neq 0$. Let

$$z = \alpha x_1 + \beta x_2 .$$

Suppose first $\alpha\beta \neq 0$. Substituting $x_2 = \frac{z - \alpha x_1}{\beta}$ in (2.1) we obtain

$$m\beta z^2 + 2ez + f\beta = 0$$

if $e\alpha - d\beta = 0$, and

$$x_1 = \frac{m\beta z_2^2 e z + f\beta}{2(e\alpha - d\beta)} \quad (3.10)$$

if $e\alpha - d\beta \neq 0$. If $e\alpha - d\beta = 0$ then z assumes one of two values, x_1 may assume any value, and (3.6) is easy to verify. If $e\alpha - d\beta \neq 0$ substitute $x_1 = \frac{z - \beta x_2}{\alpha}$ into (2.1) to get

$$x_2 = -\frac{m\alpha z^2 + 2dz + f\alpha}{2(e\alpha - d\beta)} . \quad (3.11)$$

Now the congruence class of $z \pmod{2(e\alpha - d\beta)\Gamma}$ determines whether (x_1, x_2) given by (3.10), (3.11) are integral, and if so specifies their congruence class $\pmod{\Gamma}$. Hence any specified $(x_1, x_2) \pmod{\Gamma}$ that can occur is given by some z in any block of $2(e\alpha - d\beta)\Gamma$ consecutive values of z . Now consider what signs of (x_1, x_2) can occur. The sign of x_1 changes when the numerator of the right side of (3.10) changes sign, and the sign change occurs for some z with $|z| < 6\|F\|^2$. A similar result holds for the sign of x_2 via (3.11). Hence if there is an admissible solution, there will be one with

$$|z| < 6\|F\|^2 + |2(e\alpha - d\beta)\Gamma| < 10\|F\|^3 .$$

Using (3.10) and (3.11) then gives (3.6).

Finally suppose $\alpha\beta = 0$. If $\alpha = \beta = 0$ then (2.1) is linear and (3.6) is easily verified. If $\alpha = 0$ and $\beta \neq 0$ then use (3.10) and replace (3.11) with

$$x_2 = \frac{z}{\beta} .$$

The same argument as in the case $\alpha\beta \neq 0$ now proves (3.6). The case $\alpha \neq 0, \beta = 0$ is treated similarly. $\square$

Lemma 3.2 shows we need only consider indefinite binary quadratic Diophantine equations in the sequel. In the indefinite case $ac \neq 0$ in (2.2) and hence the variable change (3.1) is invertible. We now reduce the problem of finding admissible solutions to (2.1) to that of finding an admissible solution (suitably defined) to $y_1^2 - Dy_2^2 = g$.

Lemma 3.3. *Suppose that the system (2.1)–(2.3) has an admissible solution $x = (x_1, x_2)$. Then one of the following holds.*

(i)

$$\|x\| < 200\|F\|^6 . \tag{3.12}$$

(ii) *The equation*

$$y_1^2 - Dy_2^2 = g \tag{3.13}$$

with g given by (3.3) has a solution (y_1, y_2) such that

$$y_1 > 0 \tag{3.14}$$

and one of

$$y_2 > 0 \text{ and } c(-b + \sqrt{D}) > 0 , \tag{3.15a}$$

$$y_2 < 0 \text{ and } c(b + \sqrt{D}) > 0 , \tag{3.15b}$$

holds. In addition $(y_1, y_2) \pmod{cD\Gamma}$ satisfies

$$cy_1 + c(be - cd) \equiv cD\alpha_1 \pmod{cD\Gamma} , \tag{3.16}$$

$$-by_1 + Dy_2 - c(ae - bd) \equiv cD\alpha_2 \pmod{cD\Gamma} . \tag{3.17}$$

Conversely, if the system (3.13)–(3.17) has a solution, then (2.1) has an admissible solution.

Proof. Suppose the admissible solution bx has

$$||\mathbf{x}|| \geq 200||F||^6 . \quad (3.18)$$

We show that the solution (y_1, y_2) to (3.13) given by (3.1) satisfies (3.14)–(3.17). Now (3.5) shows that (3.16), (3.17) hold.

We first show (3.18) implies $|y_1|$ and $|y_2|$ are large. If $||\mathbf{y}|| < 90||F||^5$ then absolute value estimates in (3.4) yield

$$||\mathbf{x}|| < 190||F||^6$$

contradicting (3.19). So $||\mathbf{y}|| \geq 90||F||^5$. If $|y_1| < 90||F||^5$ then $|y_2| \geq 90||F||^5$ then

$$y_1^2 \geq D(y_2)^2 - |g|$$

implies

$$|y_1| > 89||F||^5 , \quad (3.19)$$

so this holds in all cases. Then

$$y_2^2 \geq \frac{y_1^2 - |g|}{D}$$

implies

$$|y_2| \geq \frac{88}{\sqrt{D}}||F||^5 . \quad (3.20)$$

Now suppose $y_1 < 0$. Then by (3.1)

$$y_1 - (be - cd) = Dx_1 > 0$$

so

$$|y_1| \leq |be - cd| < 21||F||^2 ,$$

contradicting (3.19). Hence (3.14) holds.

To prove (3.15), note (3.13) yields

$$|y_1 + \sqrt{D} y_2| |y_1 - \sqrt{D} y_2| = |g| < 6||F||^4 . \quad (3.21)$$

Since

$$y_1 + \sqrt{D} y_2 = (y_1 - \sqrt{D} y_2) + 2\sqrt{D} y_2 ,$$

(3.20) implies that

$$MAX(|y_1 + \sqrt{D} y_2|, |y_1 - \sqrt{D} y_2|) \geq 88||F||^5 .$$

Hence (3.21) yields

$$MIN(|y_1 + \sqrt{D} y_2|, |y_1 - \sqrt{D} y_2|) < ||F||^{-1} \quad (3.22)$$

Consequently y_1 is very close to one of $\pm\sqrt{D} y_2$.

We consider first the case that

$$|y_1 - \sqrt{D} y_2| < ||F||^{-1} . \quad (3.23)$$

So $y_2 > 0$ in this case. Then (3.5) yields

$$\begin{aligned} cDx_2 &= -by_1 + Dy_2 + c(-be + cd) - eD \\ &= \sqrt{D}(-b + \sqrt{D})y_2 + \xi \end{aligned} \quad (3.24)$$

where

$$|\xi| < \|F\| + 8\|F\|^4 < 9\|F\|^4. \quad (3.25)$$

We claim

$$|-b + c\sqrt{D}| \geq \frac{1}{3}\|F\|^{-2}. \quad (3.26)$$

Indeed, suppose $|-b + \sqrt{D}| < 1$. If so, then $|-b - \sqrt{D}| > 1$. So

$$|-b + \sqrt{D}| \geq \frac{|b^2 - D|}{|-b - \sqrt{D}|} \geq \frac{1}{3\|F\|^2}.$$

Now (3.20) and (3.26) imply

$$|\sqrt{D}(-b + \sqrt{D})y_2| > 29\|F\|^4. \quad (3.27)$$

Then comparing (3.24), (3.25) and (3.27), we obtain

$$\text{sign}(cDx_2) = \text{sign}(\sqrt{D}(-b + \sqrt{D})y_2). \quad (3.28)$$

Since $x_2 \geq 0$, this yields

$$\text{sign}(y_2) = \text{sign}(c(-b + \sqrt{D}))$$

which proves (3.15a). Finally we consider the case

$$|y_1 + \sqrt{D}y_2| < \|F\|^{-2}.$$

Then $y_2 < 0$ in this case. An analysis similar to the previous case shows that

$$\text{sign}(y_2) = -\text{sign}(c(b + \sqrt{D}))$$

which gives (3.15b).

To prove the converse, suppose we have a solution (y_1, y_2) to (3.13)–(3.17) satisfying (3.15a). Let (t_1, u_1) be the minimal positive solution to Pell's equation

$$t^2 - Du^2 = 1. \quad (3.29)$$

Let

$$t_k + u_k\sqrt{D} = (t_1 + u_1\sqrt{D})^k. \quad (3.30)$$

It is well-known that (t_k, u_k) satisfy (3.29) and that for any modulus M there exists an integer $p(M)$ such that

$$\begin{aligned} t_k &\equiv 1 \pmod{M} \\ u_k &\equiv 0 \pmod{M} \end{aligned} \quad (3.31)$$

whenever

$$p(M)|k .$$

(See Appendix A.) Certainly $t_k, u_k \rightarrow \infty$ as $k \rightarrow \infty$. We now set

$$y_1^* + y_2^* \sqrt{D} = (y_1 + y_2 \sqrt{D})(t_k + u_k \sqrt{D}) \quad (3.32)$$

where $p(cd\Gamma)|k$. Note that since y_1, y_2, t_k, u_k are all positive, $y_1^* \geq t_k, y_2^* \geq u_k$. By picking k large enough, we may guarantee

$$\text{MIN}(y_1^*, y_2^*) > 88||F||^5 . \quad (3.33)$$

Furthermore (3.31) applies with $M = cd\Gamma$ guarantees that

$$y_i^* \equiv y_i \pmod{cd\Gamma} \quad i = 1, 2 . \quad (3.34)$$

Also (3.32) and (3.29) guarantee that (y_1^*, y_2^*) satisfy (3.13). Now let (x_1^*, x_2^*) be the rational solution to (2.1) associated to (y_1^*, y_2^*) by (3.5). Now (3.34) shows that (y_1^*, y_2^*) satisfies (3.16), (3.17) hence (x_1^*, x_2^*) is an integer solution and

$$x_i^* \equiv \alpha_i \pmod{\Gamma} .$$

We claim (x_1^*, x_2^*) is nonnegative. If so (x_1^*, x_2^*) is the desired admissible solution. Using (3.5) and (3.33), we obtain

$$Dx_1^* \geq y_1^* - |be - cd| > 86||F||^5$$

hence $x_1^* > 0$. Now (3.33) shows the argument (3.21)–(3.22) is valid here, and since $y_1^* > 0, y_2^* > 0$ we obtain

$$|y_1^* - \sqrt{D} y_2^*| < ||F||^{-1} .$$

The argument (3.24)–(3.28) also assumed only the truth of (3.20), so it is valid here and we obtain

$$\text{sign}(cDx_2) = \text{sign}(\sqrt{D}(-b + c\sqrt{D})y_2) .$$

We are given $y_2 > 0, c(-b + c\sqrt{D}) > 0$, hence $x_2 > 0$ follows in this case.

Now assume a solution exists to (3.13)–(3.17) satisfying (3.15b). In this case set

$$y_1^* + y_2^* \sqrt{D} = (y_1 + y_2 \sqrt{D})(t_k - u_k \sqrt{D})$$

where $p(cd\Gamma)|k$. Since $y_1 > 0, y_2 < 0, t_k > 0, u_k < 0$, we obtain $y_1^* > 0, y_2^* < 0$ and $y_1^* \geq t_k, |y_2^*| \geq |u_k|$. By picking k large enough, we ensure that

$$\text{MIN}(|y_1^*|, |y_2^*|) > 88||F||^5 . \quad (3.35)$$

Also $y_i^* \equiv y_i \pmod{cd\Gamma}$, and (y_1^*, y_2^*) satisfies (3.13). An analogous argument to the case treated above now shows that (x_1^*, x_2^*) associated to this (y_1^*, y_2^*) is an admissible solution to (2.1). $\square$

It remains to bound the size of admissible solutions to the equation

$$y_1^2 - Dy_2^2 = g . \quad (3.36)$$

The set of solutions to this equation has a simple form related to solutions of the Pell equation

$$t^2 - Du^2 = 1 .$$

Let (t_1, u_1) denote the minimal positive solution to this equation, the *fundamental solution*, and let

$$\epsilon = t_1 + u_1\sqrt{D} . \quad (3.37)$$

A solution (y_1, y_2) to (3.36) is called *basic* provided $\eta = y_1 + y_2\sqrt{D}$ has

$$1 \leq |\eta| < \epsilon . \quad (3.38)$$

Lemma 3.4. *For a positive squarefree D the complete set of solutions (x_1, x_2) to*

$$y_1^2 - Dy_2^2 = g$$

is given by

$$y_1 + y_2\sqrt{D} = \eta\epsilon^k \quad (3.39)$$

for some basic solution η and some integer k . There are only a finite number of basic solutions.

Proof. Suppose (y_1, y_2) is a solution to (3.36). Then for some integer k ,

$$\epsilon^k \leq |y_1 + y_2\sqrt{D}| < \epsilon^{k+1} .$$

Consequently for the correct choice of sign

$$\eta = \epsilon^{-k}(x_1 + x_2\sqrt{D}) \quad (3.40)$$

is a basic solution.

There are only a finite number of basic solutions since

$$\bar{\eta} \equiv y_1 - y_2\sqrt{D} = g/\eta$$

using (3.36). Hence

$$\begin{aligned} |y_1| &\leq |\eta| + |\bar{\eta}| < \epsilon + |g| \\ |y_2| &\leq \frac{\epsilon + |g|}{\sqrt{D}} . \end{aligned} \quad (3.41)$$

□

We use the following well-known upper bound on the size of the fundamental solution of Pell's equation.

Proposition 3.1. (Hua [17]) *Let (t_1, u_1) be the minimal positive solution to Pell's equation $t^2 - Du^2 = 1$. If $\epsilon = t_1 + u_1\sqrt{D}$ then*

$$\epsilon < D^{\sqrt{D}} . \quad (3.42)$$

We can now establish the following bound.

Lemma 3.5. *Suppose that the equation*

$$y_1^2 - Dy_2^2 = g \quad (3.43)$$

has an integral solution (y_1, y_2) with

$$y_i \equiv \alpha_i \pmod{M}, \quad i = 1, 2,$$

and with (y_1, y_2) having prescribed signs. Then it has such a solution with

$$\text{MAX}(\log y_1, \log y_2) \leq 9||E||^{3/2}(\log ||E||)^2 \quad (3.44)$$

where $||E|| = \text{MAX}(|D|, |g|, M)$.

Proof. Consider the set of solutions $(y_1, y_{2,k})$ to (3.43) where

$$y_{1,k} + y_{2,k}\sqrt{D} = \eta\epsilon^k, \quad (3.45)$$

k runs through the integers, and

$$\eta = y_{1,0} + y_{2,0}\sqrt{D}$$

where $(y_{1,0}, y_{2,0})$ is a fixed basic solution of (3.43). We consider these solutions from the viewpoint of their sign patterns and congruence classes $\pmod{M}$.

For sign patterns, we will show that the signs of $(y_{1,k}, y_{2,k})$ become constant for all sufficiently large positive k , and also for sufficiently large negative k . We show $\text{sign}(y_{1,k})$ is constant for all k with

$$k \geq \log |g|, \quad (3.46)$$

and is constant for all k with

$$k \leq \log |g| - 2. \quad (3.47)$$

The same holds for $y_{2,k}$. To do this, we use the standard notation $\bar{\alpha} = a - b\sqrt{D}$ for the algebraic conjugate of $\alpha = a + b\sqrt{D}$. Suppose (3.45) holds. Then

$$y_{1,k} = \frac{1}{2}(\eta(\epsilon)^k + \bar{\eta}(\bar{\epsilon})^k). \quad (3.48)$$

Now $\bar{\eta} = g/\eta$ by (3.43) and $\bar{\epsilon} = \epsilon^{-1}$. Now suppose

$$k \geq \log |g| \geq (\log \epsilon)^{-1} \log |g| \quad (3.49)$$

since the smallest ϵ that occurs is $\epsilon = 2 + \sqrt{3}$ for $D = 3$, and so $\epsilon > e$. Then since $\eta > 1$, $\bar{\epsilon}_0 < 1$,

$$|\eta(\epsilon)^k| \geq |g| > \left|\frac{g}{\eta}(\epsilon)^{-1}\right| = |\bar{\eta}(\bar{\epsilon})^k|.$$

In this case $y_{1,k}$ has the same sign as η , and is constant. Similarly when (3.47) holds we find $y_{1,k}$ has the same sign as $\bar{\eta}$. (Use the fact $1 \leq |\eta| \leq \epsilon$.) Similar arguments apply to $y_{2,k}$ using

$$y_{2,k} = \frac{1}{2\sqrt{D}}(\eta(\epsilon)^k - \bar{\eta}(\bar{\epsilon})^k).$$

For congruence conditions, in Appendix A we show that for

$$t_k + u_k \sqrt{D} = \epsilon^k \quad (3.50)$$

and any modulus M , the sequences $\{t_k\}, \{u_k\}$ formed when k varies are both periodic (mod M), and that the minimal positive period $p(M)$ of both series jointly has

$$p(M) \leq 2M(\log M + 1) . \quad (3.51)$$

Using (3.45) this guarantees that the sequences $\{y_{1,k}\}, \{y_{2,k}\}$ are both periodic (mod M) with period $p(M)$.

Combining these results, we find that all possible combinations of sign patterns and congruence conditions (mod M) that occur for $(y_{1,k}, y_{2,k})$ in the sequence (3.45) occur for some k with

$$|d| \leq 4M(\log M) + \log g + 2 . \quad (3.52)$$

In this circumstance

$$\begin{aligned} |y_{1,k}| &= \frac{1}{2} |\eta(\epsilon)^k + \bar{\eta}(\bar{\epsilon})^k| \\ &\geq \frac{1}{2} (\epsilon^{k+1} + |g|) \leq |g| \epsilon^{k+1} \end{aligned}$$

using (3.38), $|\bar{\eta}| \leq |g|$, and $\bar{\epsilon} < 1$. Hence

$$\begin{aligned} \log y_{1,k} &\leq \log g + (k+1) \log \epsilon \\ &\leq (4M(\log M) + 2 \log g + 3) \sqrt{D} \log D \end{aligned} \quad (3.53)$$

using Proposition 3.5. Using

$$|y_{2,k}| = \frac{1}{2\sqrt{D}} |\eta(\epsilon)^k - \bar{\eta}(\bar{\epsilon})^k|$$

we obtain the same bound (3.53) for $\log y_{2,k}$. The bound (3.53) implies

$$MAX(\log y_1, \log y_2) \leq 9||E||^{3/2} (\log ||E||)^2 .$$

By Lemma 3.4 all integer solutions to (3.43) fall in one of the sequences (3.45), and this proves (3.44). $\square$

Lemma 3.6. *Any indefinite binary quadratic Diophantine equation that has an admissible solution has such a solution (x_1, x_2) with*

$$MAX(\log x_1, \log x_2) \leq 210||F||^6 (\log ||F||)^2 . \quad (3.54)$$

Proof. We apply the results of Lemma 3.3 and Lemma 3.5. Using (3.35) a solution (y_1^*, y_2^*) of (3.43) will correspond to an admissible solution (x_1^*, x_2^*) of (2.1) provided (ii) of Lemma 3.3 holds and

$$\text{MIN}(|y_1^*|, |y_2^*|) \geq 88\|F\|^5 .$$

By Lemma 3.4 we may write $(y_1^*, y_2^*) = (y_{1,k}, y_{2,k})$ for some η and k in (3.45). But for

$$k \geq 7 \log \|F\| + 11$$

we have

$$|y_{1,k}| \geq \frac{1}{2}(|\eta \epsilon^k| - |\bar{\eta}(\bar{\epsilon})^k|) \tag{3.55}$$

$$\geq \frac{1}{2}[(\epsilon)^k - |g|] \tag{3.56}$$

$$\geq e^{7 \log \|F\| + 10} - |g| \geq 88\|F\|^5 . \tag{3.57}$$

A similar bound holds for $|y_{2,k}|$. We obtain the same bound for

$$k \leq -7 \log \|F\| - 13$$

using

$$\begin{aligned} |y_{1,k}| &\geq \frac{1}{2}(|\bar{\eta}(\bar{\epsilon})^k| - |\eta \epsilon^k|) \\ &\geq \frac{1}{2}[(\epsilon)^{|k|-1} - 1] . \end{aligned}$$

and similarly for $y_{2,k}$. Combining these inequalities with the argument of Lemma 3.6, we find that if (2.1) has an admissible solution (x_1, x_2) it has one whose corresponding solution $(y_{1,k}, y_{2,k})$ to (3.43) (for some η) has

$$\begin{aligned} |k| &\leq 4|cD\Gamma|(\log cD\Gamma) 10 \log \|F\| + 20 \\ &\leq 90\|F\|^4(\log \|F\|) . \end{aligned}$$

The same argument as Lemma 3.6 then gives

$$\text{MAX}(\log y_{1,k}, \log y_{2,k}) \leq 100\|F\|^5(\log \|F\|)^2$$

and (3.5) then gives (3.54). $\square$

4. Integral Binary Quadratic Forms

The problem is now reduced by Lemma 3.3 to that of studying the solutions of the equation

$$y_1^2 - Dy_2^2 = g . \tag{4.1}$$

To proceed further we use the theory of integral binary forms. A final necessary step before applying this theory is to reduce to the case where y_1, y_2 are relatively prime. If y_1, y_2 is a solution to (4.1) and $(y_1, y_2) = h$ set

$$z_1 = \frac{y_1}{h}, \quad z_2 = \frac{y_2}{h}, \quad G = \frac{g}{h^2} \quad (4.2)$$

obtaining the equation

$$\begin{aligned} z_1^2 - Dz_2^2 &= G \\ (z_1, z_2) &= 1. \end{aligned} \quad (4.3)$$

If we determine $z_1, z_2 \pmod{cD\Gamma}$ then we certainly know $y_1, y_2 \pmod{cD\Gamma}$.

An *integral binary quadratic form* $Q = [a, 2b, c]$ is given by

$$Q(x_1, x_2) = ax_1^2 + 2bx_1x_2 + c = \mathbf{x}^T \mathbf{Q} \mathbf{x}.$$

Here

$$\mathbf{x}^T = [x_1, x_2]$$

and

$$Q = \begin{bmatrix} a & b \\ b & c \end{bmatrix}$$

is a symmetric matrix *associated* to the form Q . A form Q is *properly primitive* provided $(1, 2b, c) = 1$, and we shall only deal with properly primitive forms in the rest of this paper. We say a form Q *primitively represents* an integer G provided

$$Q(z_1, z_2) = G \quad (4.4)$$

for two relatively prime integers z_1, z_2 . The *determinant* D of a form $Q = [a, 2b, c]$ is given by

$$D = b^2 - ac. \quad (4.5)$$

The *identity form* I is $[1, 0, -D]$. In this terminology (4.3) asserts that the identity form primitively represents G .

Following Gauss we will transform the question of primitive representation of integers by a form to that of determining the equivalence of two forms. A form Q_1 is *equivalent* to a form Q_2 if there is a 2×2 integer matrix $\mathbf{S}$ with $\det(\mathbf{S}) = 1$ such that

$$\mathbf{S}^t \mathbf{Q}_1 \mathbf{S} = \mathbf{Q}_2. \quad (4.6)$$

This is an equivalence relation, and we denote it by $Q_1 \approx Q_2$. This equivalence relation preserves the determinant D , the property of being a properly primitive form, and the property of primitively representing a given integer G .

Lemma 4.1. *Let z_1, z_2 satisfy (4.2). There exist z_3, z_4 such that $z_1z_4 - z_2z_3 = 1$ and the matrix $\mathbf{Z} = \begin{bmatrix} z_1 & z_3 \\ z_2 & z_4 \end{bmatrix}$ shows $I \approx Q_0$ where*

$$Q_0 = [G, 2B, C] \quad (4.7)$$

where

$$\text{MAX}(2|B|, |C|) \leq |D| + 4G^2 . \quad (4.8)$$

For any such choice of B, C, z_3, z_4 we have

$$\text{MAX}(|z_i|) \leq 6(|z_1| + |D| + G^2) . \quad (4.9)$$

$$\text{MIN}(|z_i|) \geq \frac{1}{3\sqrt{D}}(|z_1| - 5|D| - 5G^2) . \quad (4.10)$$

Proof. Choose z_3^*, z_4^* so that $z_1 z_4^* - z_2 z_3^* = 1$. Then $\mathbf{S}^* = \begin{bmatrix} z_1 & z_3^* \\ z_2 & z_4^* \end{bmatrix}$ shows $I \approx Q^*$ where

$$Q = [G, 2B^*, C^*] .$$

Now select λ so that $\mathbf{S}_1 = \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix}$ shows $Q^* \approx Q_0$ where

$$Q_0 = [G, 2B, C], \quad 0 < B < |G| .$$

Then

$$C = \frac{D - B^2}{G}$$

so

$$|C| \leq |D| + G^2 . \quad (4.11)$$

Also $I \approx Q_0$ via

$$\mathbf{Z} = \mathbf{S}^* \mathbf{S}_1 \equiv \begin{bmatrix} z_1 & z_3 \\ z_2 & z_4 \end{bmatrix} .$$

Now suppose z_3, z_4, B, C are chosen to satisfy (4.7), (4.8) and so that Z shows $I \approx Q_0$. To bound z_2, z_3, z_4 we observe that

$$G = z_1^2 - D z_2^2 \quad (4.12)$$

$$B = z_1 z_3 - D z_2 z_4 \quad (4.13)$$

$$C = z_3^2 - D z_4^2 \quad (4.14)$$

Then (4.12) gives

$$||z_1| - \sqrt{D}|z_2|| = \frac{|G|}{|z_1| + \sqrt{D}|z_2|} \leq |G| . \quad (4.15)$$

Hence

$$\frac{1}{\sqrt{D}}(|z_1| - |G|) < |z_2| < \frac{1}{\sqrt{D}}(|z_1| + |G|) . \quad (4.16)$$

Similar arguments using (4.11) and (4.4) show

$$||z_3| - \sqrt{D}|z_4|| \leq \frac{|D| G^2}{|z_3| + \sqrt{D}|z_4|} \leq |D| + G^2 \quad (4.17)$$

yielding

$$\frac{1}{\sqrt{D}}(|z_3| - |D| - G^2) \leq |z_4| \leq \frac{1}{\sqrt{D}}(|z_3| + |D| + G^2) . \quad (4.18)$$

Next note that

$$\begin{aligned}
B &= \frac{1}{2} \{ (z_1 + \sqrt{D}z_2)(z_3 - \sqrt{D}z_4) + (z_1 - \sqrt{D}z_2)(z_3 + \sqrt{D}z_4) \} \\
&= \frac{1}{2} G \left\{ \frac{z_1 + \sqrt{D}z_2}{z_3 + \sqrt{D}z_4} + \frac{z_3 + \sqrt{D}z_4}{z_1 + \sqrt{D}z_2} \right\} .
\end{aligned} \tag{4.19}$$

Viewing this as $B = \frac{1}{2}G(x + \frac{1}{x})$ then $0 < B < |G|$ gives

$$\frac{1}{3} < |x| < 3 .$$

If z_3, z_4 have the same sign, (4.9) yields

$$\frac{1}{3}(|z_1| + \sqrt{D}|z_2|) < |z_3| + \sqrt{D}|z_4| < 3(|z_1| + \sqrt{D}|z_2|) \tag{4.20}$$

so by (4.16) we obtain

$$|z_3| < 6(|z_1| + |G|) \tag{4.21}$$

and

$$|z_4| < \frac{6}{\sqrt{D}}(|z_1| + |G|) . \tag{4.22}$$

Adding (4.17) and (4.21), we obtain

$$\begin{aligned}
2|z_3| &> \frac{1}{3}(|z_1| + \sqrt{D}|z_2|) - |D| - G^2 \\
&> \frac{1}{3}(2|z_1| - 3|D| - 4G^2)
\end{aligned} \tag{4.23}$$

using (4.15). Then (4.18) gives

$$|z_4| > \frac{1}{3\sqrt{D}}(|z_1|\frac{9}{2}|D| - 5G^2) . \tag{4.24}$$

If z_3, z_4 have opposite signs, we use

$$B = \frac{1}{2} G \left\{ \frac{z_1 - \sqrt{D}z_2}{z_3 - \sqrt{D}z_4} + \frac{z_3 - \sqrt{D}z_4}{z_1 - \sqrt{D}z_2} \right\}$$

and again conclude (4.20)–(4.24) by similar arguments. $\square$

We shall simplify the situation further by replacing the form Q_0 given by Lemma 4.1 by a reduced form Q_{red} . An indefinite form $Q = [a, 2b, c]$ is *reduced* when

$$\begin{aligned}
0 &< b < \sqrt{D} \\
\sqrt{D} - b &< |a| < \sqrt{D} + b .
\end{aligned} \tag{4.25}$$

It can be checked that these inequalities imply that

$$\sqrt{D} - b < |c| < \sqrt{D} + b$$

so that

$$|Q_{\text{red}}| < 2\sqrt{D} . \quad (4.26)$$

We use the following result.

Proposition 4.1. *Given any indefinite form Q , there is a unimodular matrix $\mathbf{S}_1$ and a reduced form Q_{red} such that $Q \approx Q_{\text{red}}$ via $\mathbf{S}_1$, and*

$$\log \|\mathbf{S}_1\| = O(\log \|\mathbf{Q}\|) . \quad (4.27)$$

Proof. This is Lagarias [20, Theorem 4.1]. $\square$

The *identity reduced form*

$$\tilde{I} = [1, 2\lambda, \mu] \quad (4.28)$$

is defined by $I \approx \tilde{I}$ via

$$\mathbf{S}^* = \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix} \quad (4.29)$$

where $\lambda = [\sqrt{D}]$. $\tilde{I}$ is a reduced form. The results obtained so far are summarized in the following lemma, which will provide one part of the certificates.

Lemma 4.2. *Consider the binary quadratic Diophantine equation E given by*

$$y_1^2 - Dy_2^2 = g . \quad (4.30)$$

This equation has a solution (y_1, y_2) satisfying

$$y_i \equiv \alpha_i \pmod{M}, \quad i = 1, 2, \quad (4.31)$$

and

$$\text{sign}(y_i) = \text{sign}(\alpha_i), \quad i = 1, 2, \quad (4.32)$$

where $\text{sign}(1)$, $\text{sign}(2)$ are specified signs, provided there exist integers h, B, C and 2×2 matrices $\mathbf{S}, \mathbf{W}$ having the following properties.

(i) *h is a positive integer and $G = g/h^2$ is an integer.*

(ii) *The quadratic form $Q_0 = G, 2B, C$ is property primitive of determinant D .*

(iii) *The matrix $\mathbf{S}$ shows*

$$Q_0 \approx Q_{\text{red}} \text{ via } \mathbf{S} \quad (4.33)$$

where Q_{red} is a reduced form.

(iv) *$\mathbf{W}$ shows*

$$\tilde{I} \approx Q_{\text{red}} \text{ via } \mathbf{W} \quad (4.34)$$

where $\tilde{I}$ is the reduced identity form, Q_{red} is given by (4.33).

(v) Define $\mathbf{U} = \begin{bmatrix} u_1 & u_3 \\ u_2 & u_1 \end{bmatrix}$ by

$$\mathbf{U} = \begin{bmatrix} 1 & -\lambda \\ 0 & 1 \end{bmatrix} \mathbf{W} \mathbf{W}^{-1} \quad (4.35)$$

where $\lambda = \lceil \sqrt{D} \rceil$. The congruence class of $\mathbf{W} \pmod{M}$ is such that

$$hu_i \equiv \alpha_i \pmod{M}, \quad i = 1, 2. \quad (4.36)$$

In addition

$$\text{sign}(u_i) = \text{sign}(i), \quad i = 1, 2. \quad (4.37)$$

In fact $y_1 = hu_1, y_2 = hu_2$ then satisfy (4.30)–(4.32). Conversely, if such an admissible solution exists, then there exist integers h, B, C and 2×2 matrices $\mathbf{S}, \mathbf{W}$ having properties (i)–(v) and satisfying in addition the following bounds, where $\|E\| = \text{MAX}(D, |g|, M)$.

(vi)

$$\log h = O(\log \|E\|) \quad (4.38)$$

(vii)

$$\text{MAX}(|B|, |C|) \leq D + 4g^2, \quad (4.39)$$

(viii)

$$\log \|\mathbf{S}\| = O(\log \|E\|), \quad (4.40)$$

(ix)

$$\log \|\mathbf{W}\| = O(\|E\|^{3/2} (\log \|E\|)^2). \quad (4.41)$$

Proof. Suppose that properties (i)–(v) above hold. We check that $y_1 = hu_1, y_2 = hu_2$ satisfy (4.30)–(4.32). The congruence and sign conditions hold by (4.36), (4.37), since $h > 0$ by (i). To show (4.30) holds, we observe that

$$I \approx Q_0 \text{ via } \mathbf{U}$$

where $I = [1, 0, -D]$ and $\mathbf{U}$ is given by (4.37). For

$$(\mathbf{S}^*)^{-1} = \begin{bmatrix} 1 & -\lambda \\ 0 & 1 \end{bmatrix}$$

where $\mathbf{S}^*$ is given in (4.29) shows $I \approx \tilde{I}$, $\mathbf{W}$ shows $\tilde{I} \approx Q_{\text{red}}$ by (iv), and $\mathbf{S}^{-1}$ shows $Q_{\text{red}} \approx Q_0$ by (iii). Thus

$$\mathbf{u}^t \begin{bmatrix} 1 & 0 \\ 0 & -D \end{bmatrix} \mathbf{U} = \begin{bmatrix} G & B \\ B & C \end{bmatrix}. \quad (4.42)$$

Examining the upper left corner of this identity gives

$$u_1^2 - Du_2^2 = G.$$

Using $G = g/h^2$ by (i), (4.30) follows.

Now suppose an admissible solution (y_1, y_2) satisfying (4.30)–(4.32) exists. Using Lemma 3.5 we may suppose

$$MAX(\log y_1, \log y_2) < 9||E||^{3/2}(\log ||E||)^2 . \quad (4.43)$$

Set

$$h = g.c.d.(y_1, y_2) .$$

and $G = g/h^2$, establishing (i). Since h divides g ,

$$\log h = O(\log ||E||)$$

giving (vi). Letting $z_1 = y_1/h, z_2 = y_2/h$, we may apply Lemma 4.1 to produce B, C satisfying (ii), (vii), and a matrix $\mathbf{Z}$ showing $I \approx Q_0$. The bounds (4.8) imply

$$\log ||Q_0|| = O(\log ||E||) \quad (4.44)$$

and (4.9) gives

$$\log ||\mathbf{Z}|| = O(||E||^{3/2}(\log ||E||)^2) \quad (4.45)$$

using (4.43), since z_1 divides y_1 . Proposition 4.1 and (4.44) produces an S satisfying (iii), (viii). Take $\mathbf{U} = \mathbf{Z}$ in (4.35) and *define*

$$\mathbf{W} = \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix} \mathbf{ZS} . \quad (4.46)$$

The $u_1 = z_1, u_2 = z_2$ so (v) holds. Also $\begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix}$ shows $\tilde{I} \approx I$, $\mathbf{Z}$ shows $I \approx Q_0$ and $\mathbf{S}$ shows $Q_0 \approx Q_{\text{red}}$, hence (4.40) shows (iv) holds. Finally (4.46) gives

$$||\mathbf{W}|| \leq 8||\mathbf{S}^*|| ||\mathbf{Z}|| ||\mathbf{S}||$$

where $\mathbf{S}^* = \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix}$ with $\lambda = \lceil \sqrt{D} \rceil$. Hence

$$\log ||\mathbf{W}|| = O(||E||^{3/2}(\log ||E||)^2)$$

using (4.45) and the already established (viii). $\square$

5. Equivalence of Reduced Indefinite Binary Quadratic Forms

The problem has now been simplified to that of finding a particular matrix $\mathbf{W}$ which demonstrates the equivalence of the identity form $\tilde{I}$ and a reduced form Q_{red} . Gauss [11, Arts. 183–205] showed that such matrices have a special structure, which we describe below.

A reduced form $Q_1 = [a_1, 2b_1, c_1]$ is said to have a reduced form $Q_2 = [a_2, 2b_2, c_2]$ as a right-neighbor provided $a_2 = c_1$. In this case $Q_1 \approx Q_2$ via $\mathbf{S}$ where

$$\mathbf{S} = \begin{bmatrix} 0 & 1 \\ -1 & \lambda \end{bmatrix} \quad (5.1)$$

and λ is specified by

$$-\sqrt{D} - b_1 < \lambda c_1 < -\sqrt{D} - b_1 + |c_1| . \quad (5.2)$$

Each reduced form has a unique right neighbor. Let $Q^{(1)}$ denote the right-neighbor of $\tilde{I}$, and $Q^{(j)}$ the right-neighbor of $Q^{(j-1)}$. Let $\mathbf{S}^{(j)}$ denote the matrix given by (5.1), (5.2) taking $Q^{(j-1)}$ to $Q^{(j)}$. The set of $Q^{(j)}$ form a closed *cycle* of even *period* $2p$, i.e., there exists some $Q^{(k)} = \tilde{I}$ and the smallest $k = 2p$. We call this cycle the *principal cycle*. For $1 \leq j \leq 2p$, $\tilde{I} \approx Q^{(j)}$ via $\mathbf{L}_j$ where

$$\mathbf{L}_j = \mathbf{S}^{(1)} \dots \mathbf{S}^{(j)} . \quad (5.3)$$

We call $\mathbf{L}_j$ a *simple equivalence matrix*. The matrix $\mathbf{U} = \mathbf{L}_{2p}$ is called the *fundamental automorph*. If we set

$$\mathbf{U} = \begin{bmatrix} u & w \\ t & v \end{bmatrix}$$

the condition $\tilde{I} \approx \tilde{I}$ via $\mathbf{U}$ shows that t, u satisfies Pell's equation

$$t^2 - Du^2 = 1 . \quad (5.4)$$

In fact $(|t|, |u|)$ is the least strictly positive solution to (5.4), the *fundamental solution*, and

$$\mathbf{U} = \begin{bmatrix} u & t \\ t & Du \end{bmatrix} . \quad (5.5)$$

We may consistently extend the definition of $\mathbf{L}_j$ to apply for all integers j by first defining $\mathbf{S}^{(j)}$ for negative j by

$$\mathbf{S}^{(j)} = [\mathbf{S}^{(j_0)}]^{-1}$$

where $j \equiv j_0 \pmod{2p}$ and $0 < j_0 \leq 2p$, using (5.3) for all positive j and using

$$\mathbf{L}_{-j} = \mathbf{S}^{(-j)} \dots \mathbf{S}^{(-1)}$$

for $j > 0$. In that case, for any integer k we have

$$\mathbf{L}_{j+2kp} = \mathbf{U}^k \mathbf{L}_j .$$

Gauss proved the following result (see Mathews [25, Arts. 76, 88], Venkov [37]).

Proposition 5.1. (Gauss) *Let $\tilde{I} \approx Q$ via $\mathbf{T}$ where Q is reduced. Then there is some j with $1 \leq j \leq 2p$ such that $Q = Q^{(j)}$. Furthermore there is an integer k such that*

$$\mathbf{T} = \pm \mathbf{U}^k \mathbf{L}_j = \pm \mathbf{L}_{j+2kp} . \quad (5.6)$$

For the proof of Theorem 1.1 we need more detailed information about the equivalence matrices $\mathbf{L}_j$. We first introduce the notation that if a matrix $\mathbf{M} = [m_{ij}]$, then

$$|\mathbf{M}| = [|m_{ij}|] . \quad (5.7)$$

Lemma 5.1. *The equivalence matrices $\mathbf{L}_j$ have the following properties.*

- (i) *For $j > 0$ the entries of L_j have the sign patterns $\begin{bmatrix} ++ \\ ++ \end{bmatrix}$, $\begin{bmatrix} -+ \\ -+ \end{bmatrix}$, $\begin{bmatrix} -- \\ -- \end{bmatrix}$, $\begin{bmatrix} +- \\ +- \end{bmatrix}$ according as $j = 0, 1, 2$ or $3 \pmod{4}$.*

(ii) For $j > 0$,

$$|\mathbf{L}_j| = |\mathbf{S}^{(1)}| \dots |\mathbf{S}^{(j)}| \quad (5.8)$$

and

$$|\mathbf{L}_{-j}| = |\mathbf{S}^{(-j)}| \dots |\mathbf{S}^{(-1)}|. \quad (5.9)$$

(iii) The four entries of $\mathbf{L}_j = (l_{ij})$ are all about the same size in the sense that for any $|j| \geq 2$,

$$1 \leq \frac{\text{MAX}|l_{ij}|}{\text{MIN}|l_{ij}|} \leq 4(D + \sqrt{D}). \quad (5.10)$$

Proof. We first observe that a reduced form $Q = [a, 2b, c]$ has by definition (4.20) $|b| < \sqrt{D}$ hence

$$ac < 0. \quad (5.11)$$

The reduced forms $Q^{(i)} = [a_i, 2b_i, c_i]$ in the fundamental cycle have $a_{i+1} = c_i$. Noting $Q^{(0)} = \tilde{I}$ so $a_0 = 1$, by induction using (5.11) we obtain

$$(-1)^i a_i > 0. \quad (5.12)$$

Now (4.25) and (5.2) imply that

$$\lambda_i c_{i-1} < 0,$$

so that (5.11), (5.12) yield

$$(-1)^{i+1} \lambda_i > 0. \quad (5.13)$$

To prove (i) and (ii), note (5.13) implies $\mathbf{S}^{(i)}$ for $i > 0$ has the sign patterns $\begin{bmatrix} -+ \\ -+ \end{bmatrix}$ when i is odd, $\begin{bmatrix} ++ \\ -- \end{bmatrix}$ when i is even. Then it is easy to establish by induction on $i > 0$ that the entries of $\mathbf{L}_i$, have the sign patterns $\begin{bmatrix} ++ \\ ++ \end{bmatrix}$, $\begin{bmatrix} -+ \\ -+ \end{bmatrix}$, $\begin{bmatrix} -- \\ -- \end{bmatrix}$, $\begin{bmatrix} +- \\ +- \end{bmatrix}$ according as $i = 0, 1, 2, 3 \pmod{4}$. Another induction on $i > 0$ shows no cancellation occurs in multiplying the entries of $\mathbf{L}_i$ by $\mathbf{S}^{(i+1)}$ and (5.8) follows. For $i < 0$, we observe first that

$$\begin{bmatrix} 0 & l \\ -1 & \lambda \end{bmatrix}^{-1} = \begin{bmatrix} \lambda & -1 \\ 1 & 0 \end{bmatrix}.$$

Then note $\lambda_i = \lambda_{i-2p}$ so (5.13) holds for $i < 0$ as well. This implies that for $i < 0$, $\mathbf{S}^{(i)}$ has the sign patterns $\begin{bmatrix} -- \\ ++ \end{bmatrix}$ if i is odd, $\begin{bmatrix} +- \\ +- \end{bmatrix}$ if i is even. Another induction shows for $i < 0$ that the entries of $\mathbf{L}_i$ have the sign patterns $\begin{bmatrix} ++ \\ ++ \end{bmatrix}$, $\begin{bmatrix} -- \\ ++ \end{bmatrix}$, $\begin{bmatrix} -- \\ -- \end{bmatrix}$, $\begin{bmatrix} ++ \\ -- \end{bmatrix}$ according as $i \equiv 0, 1, 2$ or $3 \pmod{4}$. Then (5.9) follows by induction.

To prove (iii), consider first the case $j > 0$. Using (5.8), we need only bound the entries

$$|\mathbf{L}_j| = \left| \begin{array}{cc} 0 & 1 \\ 1 & |\lambda_1| \end{array} \right| \cdots \left| \begin{array}{cc} 0 & 1 \\ 1 & |\lambda_j| \end{array} \right|. \quad (5.14)$$

The formulae for $|\mathbf{L}_j|$ is exactly that of the ordinary continued fraction algorithm, where

$$|\mathbf{L}_j| = \begin{vmatrix} p_{j-1} & p_j \\ q_{j-1} & q_j \end{vmatrix} \quad (5.15)$$

and $\frac{p_j}{q_j}$ is the j th convergent to $\theta = [0, |\lambda_1|, |\lambda_2|, \dots]$. In particular, for any $j \geq 2$ we have

$$\frac{|\lambda_2|}{|\lambda_1\lambda_2| + 1} = \frac{p_3}{q_3} \leq \frac{p_j}{q_j} \leq \frac{p_2}{q_2} = \frac{1}{|\lambda_1|} . \quad (5.16)$$

Now (5.2) implies

$$1 \leq |\lambda_i| < 2\sqrt{D} \quad (5.17)$$

so (5.15) yields

$$\frac{1}{2\sqrt{D} + 1} \leq \frac{p_j}{q_j} < 1 . \quad (5.18)$$

In addition

$$q_{j+1} = |\lambda_j|q_j + q_{j-1} \leq (|\lambda_j| + 1)q_j . \quad (5.19)$$

Combining (5.16), (5.17), we obtain

$$\begin{aligned} p_j &\leq p_{j+1} \leq q_{j+1} \\ p_j &\leq q_j \leq q_{j+1} . \end{aligned} \quad (5.20)$$

Finally

$$q_{j+1} \leq (2\sqrt{D} + 1)q_j \leq (2\sqrt{D} + 1)^2 p_j \quad (5.21)$$

and since $p_j \geq 1$ for $j \geq 2$ by (5.15) this implies (5.10) on this range. The case $j < 0$ is treated analogously to $j > 0$. In this case

$$|\mathbf{L}_{-j}| = \begin{vmatrix} q_{j+1} & q_j \\ p_{j+1} & p_j \end{vmatrix} ,$$

however. $\square$

We remark in passing that there is a close connection between the λ_i and the continued fraction expansion of $\sqrt{D}$. Let

$$\sqrt{D} = [\mu_0, \overline{\mu_1, \dots, \mu_n}]$$

where $[\mu_1, \dots, \mu_n]$ is the purely periodic part of the expansion and n is the shortest period. Then $n = 2p$ if n is even, and $n = p$ otherwise. In either case $\mu_i = |\lambda_i|$ for $1 \leq i \leq 2p$.

Our next step is to estimate the size of the entries of $\mathbf{L}_j$ in relation to j .

Lemma 5.2. *For all $j > 0$,*

$$\log \|\mathbf{L}_{j+2}\| \geq \log \|\mathbf{L}_j\| + 1 , \quad (5.22)$$

and

$$\log \|\mathbf{L}_j\| \leq \log \|\mathbf{L}_{j+1}\| \leq \log \|\mathbf{L}_j\| + \log D + 2 . \quad (5.23)$$

Proof. For the bound (5.22), we use (5.14), (5.15) to obtain

$$q_{j+2} = (|\lambda_{j+1}\lambda_j| + 1)q_j + |\lambda_{j+1}|q_{j-1} \geq 2q_j .$$

The left side of (5.23) follows from (5.21). Finally (5.21) implies

$$\log \|\mathbf{L}_{j+1}\| \leq \log \|\mathbf{L}_j\| + \log(2\sqrt{D} + 1)$$

from which the right side of (5.23) follows.

Analogous inequalities to (5.22), (5.23) hold for $j < 0$. Using Lemma 5.2 it is easy to prove by induction that for $j \geq 1$ we have

$$\frac{1}{2}|j| \leq \log \|\mathbf{L}_j\| \leq |j|(\log D + 2) . \quad (5.24)$$

The same holds for $j \leq -1$. $\square$

We next state a bound for the length $2p$ of the fundamental cycle.

Proposition 5.2. *The period $2p$ of the fundamental cycle satisfies*

$$p < (\sqrt{D} + 1) \log D . \quad (5.25)$$

Proof. The result of Hua [17] asserts that if (t_0, u_0) is the fundamental positive solution to $x^2 - Dy^2 = 1$ then

$$\frac{t_0 + u_0\sqrt{D}}{2} < D^{\sqrt{D}} .$$

Using (5.5) we obtain

$$\|\mathbf{L}_{2p}\| \leq D^{\sqrt{D}+1} . \quad (5.26)$$

Combining (5.26) with (5.24) gives

$$p \leq \log \|\mathbf{L}_{2p}\| \leq (\sqrt{D} + 1) \log D . \quad \square$$

The examples $D = 5^{2n+1}$ mentioned in the introduction show that periods $p > \frac{1}{c}\sqrt{D}$ do occur.

We can use the preceding results determine properties of the matrix $\mathbf{W}$ of Lemma 4.2.

Lemma 5.3. *Let $\mathbf{W}$ be the matrix guaranteed to exist in Lemma 4.2, satisfying (i)–(ix) of that lemma. Then*

$$\mathbf{W} = \pm(\mathbf{L}_{2p})^k \mathbf{L}_j \quad (5.27)$$

for some j with $1 \leq j \leq 2p$ and an integer k satisfying

$$|k| = O(\|E\|^{3/2}(\log \|E\|)^2) . \quad (5.28)$$

Proof. This follows immediately from Proposition 5.1, Lemma 5.2 and (4.41). $\square$

6. Composition of Binary Quadratic Forms

The certificates we construct use the operation of composition of binary quadratic forms introduced by Gauss, in particular the action of composition on the fundamental cycle of reduced forms. The idea of analyzing the action of composition on the cycle of reduced forms equivalent to $\tilde{I}$ is due to D. Shanks [33].

The simplest example of composition of two binary quadratic forms is the identity

$$(x_1^2 + x_2^2)(y_1^2 + y_2^2) = (x_1y_1 - x_2y_2)^2 + (x_1y_2 + x_2y_1)^2 \quad (6.1)$$

noted by Fermat, which shows that the product of two numbers which are the sum of two squares is itself the sum of two squares. We can rewrite (6.1) in the form

$$Q_1(x_1, x_2)Q_1(y_1, y_2) = Q_1(x_1y_1 - x_2y_2, x_1y_2 + x_2y_1) \quad (6.2)$$

where $Q_1 = [1, 0, 1]$, and in matrix terms as

$$\mathbf{x}^t \mathbf{Q}_1 \mathbf{x} \mathbf{y}^t \mathbf{Q}_1 \mathbf{y} = \mathbf{z}^t \mathbf{B}^t \mathbf{Q}_1 \mathbf{B} \mathbf{z} \quad (6.3)$$

where

$$\begin{aligned} \mathbf{x}^t &= [x_1, x_2], & \mathbf{y}^t &= [y_1, y_2] \\ \mathbf{z}^t &= [x_1y_1, x_1y_2, x_2y_1, x_2y_2] \end{aligned} \quad (6.4)$$

and

$$\mathbf{B} = \begin{bmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 1 & 0 \end{bmatrix}. \quad (6.5)$$

In this case we say Q_1 is composed of Q_1 and Q_1 via the bilinear matrix $\mathbf{B}$ of (6.5).

In the general case we say a quadratic form $Q_3 = [a_3, 2b_c, c_3]$ is *composed* of forms $Q_1 = [a_1, 2b_1, c_1]$ and $Q_2 = [a_2, 2b_2, c_2]$ via a *bilinear matrix* $\mathbf{B}$ provided the matrix equation

$$\mathbf{x}^t \mathbf{Q}_1 \mathbf{x} \mathbf{y}^t \mathbf{Q}_2 \mathbf{y} = \mathbf{z}^t \mathbf{B}^t \mathbf{Q}_3 \mathbf{B} \mathbf{z} \quad (6.6)$$

holds, where $\mathbf{x}, \mathbf{y}, \mathbf{z}$ are given by (6.4), the x_i and y_j are indeterminates, and $\mathbf{B}$ is a 2×4 bilinear matrix required to be *unimodular* and *oriented* (terms defined below). A 2×4 matrix $\mathbf{B} = [b_{ij}]$ is:

(i) *unimodular* provided the six cofactors

$$\Delta_{ij} = \begin{bmatrix} b_{1i} & b_{1j} \\ b_{2i} & b_{2j} \end{bmatrix}, \quad 1 \leq i < j \leq 4$$

have greatest common divisor 1.

(ii) *oriented* provided $a_1 \Delta_{12} > 0$ and $a_2 \Delta_{13} > 0$.

We write $Q_3 = Q_1 \circ Q_2$ to indicate composition of forms.

Our treatment of composition of forms is based on Mathews [25], Venkov [37] and Lagarias [20]. One may also consult Buell [8], Gauss [15], Shanks [32], Smith [35]. If Q_3 is composed of two properly primitive forms Q_1 and Q_2 of determinant D , then Q_3 itself is properly primitive of determinant D .

In the rest of this section we deal only with *properly primitive* indefinite forms. We first recall the following result on composition.

Proposition 6.1. *Given any two reduced forms Q_1, Q_2 of determinant D there is a reduced form Q_3 of determinant D and a bilinear matrix $\mathbf{B}$ such that $Q_3 = Q_1 \circ Q_2$ via $\mathbf{B}$ and*

$$\log \|\mathbf{B}\| = O(\log D) . \quad (6.7)$$

Proof. This is shown in Lagarias [20, Theorem 5.5]. $\square$

The key result facilitating the use of composition to create short certificates is the following lemma. Before stating it, we recall that the Kronecker product $\mathbf{S} \otimes \mathbf{T}$ of an $m \times n$ matrix $\mathbf{S} = [s_{ij}]$ and a $k \times l$ matrix $\mathbf{G}$ is a $km \times ln$ matrix

$$S \otimes T = \begin{bmatrix} s_{11}T & \cdots & s_{1n}T \\ \vdots & & \vdots \\ s_{m1}T & \cdots & s_{mn}T \end{bmatrix}$$

given in block matrix form.

Lemma 6.1. *Let $\tilde{I} \approx Q_1$ via $\mathbf{S}_1$ and $\tilde{I} \approx Q_2$ via $\mathbf{S}_2$. If $Q_3 = Q_1 \circ Q_2$ via $\mathbf{B}$, then $\tilde{I} \approx Q_3$ via $\mathbf{S}_3$ where $\mathbf{S}_3$ satisfies the matrix equation*

$$\mathbf{S}_3 \mathbf{B} = \mathbf{B}_0 (\mathbf{S}_1 \otimes \mathbf{S}_2) \quad (6.8)$$

where

$$\mathbf{B}_0 = \begin{bmatrix} 1 & 0 & 0 & D - \lambda^2 \\ 0 & 1 & 1 & 2\lambda \end{bmatrix} \quad (6.9)$$

and $\lambda = [\sqrt{D}]$.

Proof. It is straightforward to check $I = I \circ I$ via

$$\mathbf{B} = \begin{bmatrix} 1 & 0 & 0 & D \\ 0 & 1 & 1 & 0 \end{bmatrix} .$$

Using Lagarias [20, Lemma 5.1 (i)], since $I \approx \tilde{I}$ via $\begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix}$ we obtain $\tilde{I} = \tilde{I} \circ \tilde{I}$ via $\mathbf{B}_0$. using the same [20, Lemma 5.1 (i)], we next conclude $\tilde{I} = Q_1 \circ Q_2$ via $\mathbf{B}_0(\mathbf{S}_1 \oplus \mathbf{S}_2)$. Then using [20, Lemma 5.1 (ii)], we conclude there exists a unimodular integer matrix $\mathbf{S}_3$ such that

$$\mathbf{S}_3 \mathbf{B} = \mathbf{B}_0 (\mathbf{S}_1 \otimes \mathbf{S}_2) ,$$

the desired result. $\square$

We note that $\mathbf{S}_3$ is uniquely determined by (6.8), since $\mathbf{B}$ contains an invertible 2×2 submatrix by the unimodularity condition.

Now suppose Q_1 and Q_2 are forms in the principal cycle. Lemma 6.1 shows that if $Q_3 = Q_1 \circ Q_2$ and Q_3 is reduced, then Q_3 is also in the principal cycle. By Proposition 5.1 there are integers k_1, k_2 and k_3 such that $\mathbf{S}_i = \pm \mathbf{L}_{k_i}$ for $1 \leq i \leq 3$. What is the relation among the k_i 's?

Lemma 6.2. *Let Q_1, Q_2, Q_3 be in the principal cycle and suppose $\tilde{I} \approx Q_1$ via $\pm \mathbf{L}_{k_1}$, $\tilde{I} \approx Q_2$ via $\pm \mathbf{L}_{k_2}$, where $k_1, k_2 \geq 0$. Suppose $Q_3 = Q_1 \circ Q_2$ via $\mathbf{B}$ and that*

$$\log \|\mathbf{B}\| \leq c_1 \log D . \quad (6.10)$$

Let $\mathbf{S}_3$ be defined by

$$\mathbf{S}_3 \mathbf{B} = \mathbf{B}_0 (\mathbf{S}_0 (\mathbf{S}_1 \otimes \mathbf{S}_2)) .$$

If $\mathbf{S}_i = \pm \mathbf{L}_{k_i}$ and ξ is defined by

$$\log \|\mathbf{L}_{k_3}\| = \log \|\mathbf{L}_{k_1}\| + \log \|\mathbf{L}_{k_2}\| + \xi ,$$

then we have

$$|\xi| \leq (c_1 + 4) \log D . \quad (6.11)$$

Proof. By (6.8) we have

$$\|\mathbf{S}_3 \mathbf{B}\| = \|\mathbf{B}_0 (\mathbf{S}_1 \otimes \mathbf{S}_2)\| . \quad (6.12)$$

Now

$$\begin{aligned} \|\mathbf{S}_0 (\mathbf{S}_1 \otimes \mathbf{S}_2)\| &\leq 4 \|\mathbf{B}_0\| \|\mathbf{S}_1 \otimes \mathbf{S}_2\| \\ &= 4 \|\mathbf{B}_0\| \|\mathbf{S}_1\| \|\mathbf{S}_2\| . \end{aligned} \quad (6.13)$$

We next note that $\mathbf{B}_0$ is nonnegative and that $\mathbf{S}_1 \otimes \mathbf{S}_2$ has constant sign on columns by Lemma 5.2 (i). This implies

$$\|\mathbf{B}_0 (\mathbf{S}_1 \otimes \mathbf{S}_2)\| \geq \|\mathbf{S}_1 \otimes \mathbf{S}_2\| = \|\mathbf{S}_1\| \|\mathbf{S}_2\| . \quad (6.14)$$

On the other hand

$$\|\mathbf{S}_3 \mathbf{B}\| \leq 2 \|\mathbf{B}\| \|\mathbf{S}_3\| . \quad (6.15)$$

Using orientability the first two columns of $\mathbf{B}$ form an invertible 2×2 submatrix $\mathbf{B}_1$ and we obtain

$$\|\mathbf{S}_3 \mathbf{B}\| \geq \|\mathbf{S}_3 \mathbf{B}_1\| \geq \frac{\|\mathbf{S}_3\|}{2 \|\mathbf{B}_1\|} \geq \frac{\|\mathbf{S}_3\|}{2 \|\mathbf{B}_1\|} \geq \frac{\|\mathbf{S}_3\|}{2 \|\mathbf{B}\|} , \quad (6.16)$$

where the center inequality is deduced from

$$\|\mathbf{S}_3\| \leq 2 \|\mathbf{S}_3 \mathbf{B}_1\| \|\mathbf{B}_1^{-1}\|$$

and

$$\|\mathbf{B}_1^{-1}\| = (\det B)^{-1} \|\mathbf{B}\| \leq \|\mathbf{B}\| .$$

Now (6.12), (6.14), (6.15) yield

$$\|\mathbf{S}_3\| \geq \frac{2}{\|\mathbf{B}\|} (\|\mathbf{S}_1\| \|\mathbf{S}_2\|) , \quad (6.17)$$

while (6.12), (6.13), (6.16) yield

$$\|\mathbf{S}_3\| \leq 8 \|\mathbf{B}_0\| \|\mathbf{B}\| (\|\mathbf{S}_1\| \|\mathbf{S}_2\|) . \quad (6.18)$$

Using

$$\log \|\mathbf{B}_0\| \leq \log D$$

and the hypothesis (6.10), the inequalities (6.17) and (6.18) establish (??). $\square$

Lemma 5.2 shows that, for $j \geq 1$, $\log \|\mathbf{L}_j\|$ provides a measure of the size of the subscript j . On the other hand, (??) shows that composing a form $Q^{(j)}$ with itself essentially doubles this size. By repeatedly doubling the size we can rapidly move to forms far apart in the principal cycle.

Lemma 6.3. *For any $\mathbf{L}_j$ with $1 \leq j \leq 2p$ there is a sequence of equivalence matrices $\mathbf{V}_k$, and reduced forms $\tilde{Q}_k$ of length K such that*

$$(i) \quad \tilde{Q}_0 = \tilde{I}, \mathbf{V}_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

(ii) *Each pair $(\tilde{Q}_{k+1}, \mathbf{V}_{k+1})$ is obtained from the preceding $(\tilde{Q}_k, \mathbf{V}_k)$ by a transformation of either Type I or Type II, where:*

Type I. $\tilde{Q}_{k+1}$ is the right-neighbor of $\tilde{Q}_k$ so that

$$\tilde{\mathbf{Q}}_{k+1} = \mathbf{S}_{k+1}^t \tilde{\mathbf{Q}}_k \mathbf{S}_{k+1}, \quad (6.19)$$

$$\mathbf{V}_{k+1} = \mathbf{V}_k \mathbf{S}_{k+1} \quad (6.20)$$

and

$$\log \|\mathbf{S}_k\| \leq \frac{1}{2}(\log D). \quad (6.21)$$

Type II. $\tilde{Q}_{k+1} = \tilde{Q}_{k_1} \circ \tilde{Q}_{k_2}$ via $\mathbf{B}_{k+1}$ for some $0 \leq k_1, k_2 \leq k$ so that

$$\mathbf{x}^t \tilde{\mathbf{Q}}_i \mathbf{x} \mathbf{y}^t \tilde{\mathbf{Q}}_j \mathbf{y} = \mathbf{z}^t \mathbf{B}_{k+1}^t \tilde{\mathbf{Q}}_{k+1} \mathbf{B}_{k+1} \mathbf{z}, \quad (6.22)$$

$$\mathbf{V}_{k+1} \mathbf{B}_{k+1} = \mathbf{B}_0 (\mathbf{V}_{k_1} \otimes \mathbf{V}_{k_2}), \quad (6.23)$$

and where

$$\log \|\mathbf{B}_{k+1}\| = O(\log D). \quad (6.24)$$

(iii)

$$Q_K = Q^{(j)} \text{ and } \mathbf{V}_K = \mathbf{L}_j. \quad (6.25)$$

The length K of this sequence satisfies

$$K = 9(\log D)^2. \quad (6.26)$$

Proof. We suppose that the composition of reduced forms is done as in Proposition 6.1, so that (6.24) is satisfied. We let c_1 denote the constant implied by the O-symbol in (6.24). Let σ_j denote the minimal number of type I and type II transformations sequentially applied to get from $\tilde{I}$ to $Q^{(j)}$ via $\mathbf{L}_j$. First note

$$\sigma_j \leq j \quad (6.27)$$

by using type I transformations only. We will prove by induction on j that for

$$2(c_1 + 6) \log D \leq j \leq 2p \quad (6.28)$$

we have

$$\sigma_j \leq (5 + 4(c_1 + 4) \log D)(\log \|\mathbf{L}_j\|) . \quad (6.29)$$

Suppose (6.28) holds. Take j_1 to be some l such that

$$-(c_1 + 5) \log D - 2 < \log \|\mathbf{L}_l\| - \frac{1}{2} \log \|\mathbf{L}_j\| \leq -(c_1 + 4) \log D . \quad (6.30)$$

At least one such l exists by (5.23) and $1 \leq l < j$. (Note (5.24) shows $\frac{1}{2} \log \|\mathbf{L}_j\| - (c_1 + 4) \log D \geq 2$.) Hence we can obtain $\tilde{Q}_k = Q^{(j_1)}$, $\mathbf{V}_k = \mathbf{L}_{j_1}$ where $k = \sigma_{j_1}$ satisfies (6.29) by the induction hypothesis. Now apply a type II transformation, using $\tilde{Q}_k \circ \tilde{Q}_k$, obtaining $\tilde{Q}_{k+1} = Q^{j_2}$ and $\mathbf{V}_{k+1} = \mathbf{L}_{j_2}$. Using Lemma 6.3 and (6.30) we have

$$2(c_1 + 4) \log D + 2 \geq \log \|\mathbf{L}_j\| - \log \|\mathbf{L}_{j_2}\| \geq 0 .$$

Then Lemma 5.3 implies

$$0 \leq j - j_2 < 4(1 + (c_1 + 4) \log D) .$$

Hence $4(1 + (c_1 + 4) \log D)$ type I transformations will take us to $Q^{(j)}$, $\mathbf{L}_j$. Hence

$$\sigma_j \leq \sigma_{j_1} + 4((c_1 + 4) \log D) + 5 . \quad (6.31)$$

But the right side inequality of (6.30) gives

$$\log \|\mathbf{L}_{j_1}\| \leq \frac{1}{2} \log \|\mathbf{L}_j\| \leq \log \|\mathbf{L}_j\| - 1 . \quad (6.32)$$

Substituting (6.29) for j_1 into (6.31) and using (6.32) establishes (6.29) for j and completes the induction step. $\square$

Remark. By a more complicated argument, this bound (6.26) can be improved to

$$K = O(\log D) .$$

7. Certificates for Equivalence of Two Indefinite Binary Quadratic Forms

Lemma 6.3 can immediately be used to provide certificates for the equivalence of two indefinite binary quadratic forms.

Theorem 7.1. *Let Q_1 and Q_2 be two indefinite binary quadratic forms with the same discriminant. If $Q_1 Q_2$, there is a certificate of this equivalence requiring at most*

$$O(\log \|Q_1\| + \log \|Q_2\| + (\log D)^2 M(\log D)) \quad (7.1)$$

elementary operations to verify.

Proof. A necessary condition for the equivalence of two forms $Q_1 = [a_1, 2b_1c]$ and $Q_2 = [a_2, 2b_2, c_2]$ is that

$$G.C.D.(a_1, b_1, c_1) = G.C.D.(a_2, b_2, c_2) = \sigma_1$$

and

$$G.C.D.(a_1, 2b_1, c_2) = G.C.D.(a_2, 2b_2, c_2) = \sigma_2 .$$

By removing σ_1 from the coefficients of both Q_1 and Q_2 we need only consider the case $\sigma_1 = 1$. In that case the forms are *properly primitive* if $\sigma_2 = 1$ and *improperly primitive* if $\sigma_2 = 2$.

Suppose first the forms are properly primitive. Replace $Q_2 = [a_2, 2b_2, c_2]$ by $\bar{Q}_2 = [a_2 - 2b_2, c_2]$, its *inverse* form. Reduce Q_1 and $\bar{Q}_2$, obtaining $Q_1^*, \bar{Q}_2^*$. This requires $O(\log ||Q_1|| + \log ||Q_2||)$ operations by Proposition 4.1. Compose Q_1^* and $\bar{Q}_2^*$ to obtain a reduced form Q_3^* . By Proposition 6.1 this can be done in $9(M(\log D))$ operations.

Now $Q_1 \approx Q_2$, if and only if $Q_3^* \approx \tilde{I}$. This follows from the well-known facts that composition of forms induces the structure of an abelian group on equivalence classes $[Q]$ of properly primitive forms Q , that $[\tilde{I}]$ is the identity element of this group, and that $[Q]^{-1} = [\bar{Q}]$. (e.g. see Mathews [25, Arts. 141, 145].)

We now take the sequence of reduced forms $\tilde{Q}_k$ showing $Q_3^* \approx \tilde{I}$ that are guaranteed to exist by Lemma 6.3, together with the matrices $\mathbf{S}_k$ and $\mathbf{B}_k$ involved in the corresponding type I or II transformation. For each transformation we verify either (6.19) or (6.22), and this requires $O(M(\log D))$ elementary operations. We obtain a total of $O((\log D)^2 M(\log D))$ elementary operations in all, by (6.26).

Finally, we verify by induction on k that checking (6.19), (6.22) at each step guarantees that all $\tilde{Q}_k = \tilde{I}$. Certainly $\tilde{Q}_0 \approx \tilde{I}$. If a type I transformation is used, then $\tilde{Q}_{k+1} \approx \tilde{Q}_k \approx I$ by definition of equivalence. If a type II transformation is used, then $\tilde{Q}_i \approx \tilde{I}$ and $\tilde{Q}_j \approx \tilde{I}$ guarantees $\tilde{Q}_{k+1} = \tilde{Q}_1 \circ \tilde{Q}_j \approx I$ by Lemma 6.1. This completes the proof in the properly primitive case.

We take care of the improperly primitive case by reducing it to the properly primitive case by the following expedient. (Mathews [25, Art. 153].) We first note that improperly primitive forms have $D \equiv 1 \pmod{4}$. Let

$$Q \left| \begin{bmatrix} a & b \\ c & d \end{bmatrix} \right| \equiv Q(ax + by, cx + dy) .$$

If $D \equiv 1 \pmod{8}$ and Q is improperly primitive, then $Q \left| \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \right| = 2Q^*$ where Q is properly primitive. Furthermore if Q_1, Q_2 are two such improperly primitive forms then $Q_1 \approx Q_2$ if and only if $Q_1^* \approx Q_2^*$. We may find a certificate for this as above. If $D \equiv 5 \pmod{8}$ and Q is improperly primitive, then $Q \left| \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \right| = 2Q^{(1)}$, $Q \left| \begin{bmatrix} 0 & 1 \\ 0 & 2 \end{bmatrix} \right| = 2Q^{(2)}$ and $Q \left| \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} \right| = 2Q^{(i)}$ where the $Q^{(i)}$ are all properly primitive. Furthermore if Q_1, Q_2 are two such improperly primitive forms then $Q_1 \approx Q_2$ if and only if one of $Q_1^{(i)} \approx Q_2^{(1)}$ for $1 \leq i \leq 3$. We may find a certificate for this as above. In order to get the bound (7.1) we first reduce the improperly primitive forms and then apply the procedure above. This reduction involves only $O(\log D)$ additional operations.

□

Remark. Since $\|Q\| > \frac{1}{2}\sqrt{D}$ for any form Q , (7.1) gives a bound polynomial in the length of the input $\log \|Q_1\| + \log \|Q_2\|$.

8. Succinct Certificates

We now prove the main result.

Proof of Theorem 1.1. If (2.1) has an admissible solution with $\|\mathbf{x}\| < 256\|F\|^8$, then it serves as the certificate, and only $O(M(\log \|F\|))$ operations are needed to verify it is one. By Lemma 3.2 this is always the case for definite or degenerate binary quadratic Diophantine equations.

Now suppose (2.1) is indefinite, and has admissible solutions, but none with $\|\mathbf{x}\| < 256\|F\|^8$. Then by part (ii) of Lemma 3.3, there exists β_1, β_2 such that the

$$y_1^2 - Dy_2^2 = g. \quad (8.1)$$

has a solution with

$$y_i \equiv \beta_i \pmod{cD\Gamma} \quad (8.2)$$

satisfying (3.16), (3.17), and $y_1 > 0$ and the sign of y_2 is specified and satisfies one of (3.15a), (3.15b). Call the system (8.1), (8.2) with the given sign conditions E , and observe that

$$\|E\| \leq \text{MAX}(D, |g|, |cD\Gamma|) \leq 6\|F\|^4 \quad (8.3)$$

using (3.4). By Lemma 3.3 it suffices to give a certificate for this equation, to guarantee (2.1) has an admissible solution. Note that it takes only $O(M(\log D) \log D)$ operations to check the conditions of (ii) of Lemma 3.3 hold, in particular $O(M(\log D) \log D)$ operations to compute $\sqrt{D}$ to one digit past the decimal point in (3.15). Note $\log \|E\| = 9(\log \|F\|)$.

Lemma 4.2 shows that to show the system E has an admissible solution it suffices to produce certificates showing there exist integers h, B, C and 2×2 matrices $\mathbf{S}, \mathbf{W}$ such that (i)–(v) of that lemma hold. The rest of the proof will accomplish this.

Lemma 4.2 also shows that there exist integers h, B, C and 2×2 matrices $\mathbf{S}, \mathbf{W}$ such that (i)–(ix) of that lemma hold. In the rest of the proof we shall fix this particular choice of $h, B, C, \mathbf{S}$, and $\mathbf{W}$, as well as

$$Q_{\text{red}} = [a_0, 2b_0, c_0] \quad (8.4)$$

arising in (iii) of that lemma. In that case (i), (ii) of Lemma 4.2 can be verified in $O(M(\log \|E\|))$ operations by (vi) of that Lemma and (8.3). To verify (iii) of Lemma 4.2 we note that it asserts that

$$\begin{bmatrix} a_0 & b_0 \\ b_0 & c_0 \end{bmatrix} = \mathbf{S}^t \begin{bmatrix} G & B \\ B & C \end{bmatrix} \mathbf{S}. \quad (8.5)$$

Using the bound (4.26) for a reduced form, (vii), (viii) of Lemma 4.2, and (8.3), all entries in (8.5) are $O(\log \|E\|)$ so (8.5) can be verified in $O(M(\log \|E\|))$ operations.

The essential difficulty in producing the certificates is the possible large size of the entries of $\mathbf{W}$, evidenced by the bound (4.41), so that we cannot afford to keep track of these entries as fixed point binary integers. Consequently (iv) and (v) of Lemma 4.2 must be verified indirectly.

The certificates verifying (iv) and (v) are based on two kinds of formulae, which we call *short* and *long*. The short formulae can be evaluated using fixed-point integer arithmetic. We will use these to verify (iv). The long formulae involve integers with too many binary digits to allow direct evaluation. We use these to verify (v), by evaluating them (mod $cD\Gamma$) to verify (4.36), and by evaluating them using floating-point arithmetic to enough accuracy to verify (4.37).

The formulae are those guaranteed to exist by Lemma 5.3 and Lemma 6.3. By Lemma 5.3 the $\mathbf{W}$ of Lemma 4.2 can be written in the form

$$\mathbf{W} = (-1)^m (\mathbf{L}_{2p})^k \mathbf{L}_j \quad (8.6)$$

for some j with $1 \leq j \leq 2p$, for some $m = 0$ or 1 , and for which

$$k = O(\|E\|^{17} \log \|E\|) . \quad (8.7)$$

Equation (8.6) is a long formula.

Next, by Lemma 6.3 for each $\mathbf{L}_{2p}$ and $\mathbf{L}_j$ there exists a chain of reduced forms $\tilde{Q}_k$ and equivalence matrices $\mathbf{V}_k$ having the properties (6.19)–(6.26). Formulae (6.19) and (6.22) are short formulae, while (6.20) and (6.23) are long formulae.

Consider the short formulae for $\mathbf{L}_j$ and $\mathbf{L}_{2p}$. The bounds (4.26), (6.21) and (6.24) imply that each formula can be evaluated exactly using fixed-point integer arithmetic with $O(\log \|E\|)$ binary digits. Each evaluation takes $O(M(\log \|E\|))$ operations, so (6.26) implies a total of $O(M(\log \|E\|)(\log \|E\|)^2)$ operations at most in evaluating all the short formulae. In addition we must check that the bilinear matrices $\mathbf{B}_i$ used in short formulae are unimodular and oriented. Using the Euclidean algorithm to check unimodularity takes $O(M(\log \|E\|) \log \|E\|)$ operations for each $\mathbf{B}_k$, by [20, Prop. 3.3], for a total of $O(M(\log \|E\|)(\log \|E\|)^3)$ operations in all. Checking orientability requires $O(M(\log \|E\|) \log \|E\|)$ operations in all.

We now verify (iv) of Lemma 4.2. Since $\tilde{I} \approx \tilde{I}$ via $\mathbf{L}_{2p}$ and $\tilde{I} \approx Q^{(j)}$ via $\mathbf{L}_j$, (8.16) implies that

$$\tilde{I} \approx Q^{(j)} \text{ via } \mathbf{W} .$$

In order to verify (iv) it suffices to check that

$$Q^{(j)} = Q_{\text{red}} \quad (8.8)$$

where Q_{red} is as in (4.33), $Q^{(j)}$ is the Q_k produced in (6.25) for $\mathbf{L}_j$. Checking (8.8) takes another $O(\log \|E\|)$ operations.

We now describe certificates for (v) of Lemma 4.2. We first must verify

$$hu_i \equiv \alpha_i \pmod{cD\Gamma} \quad i = 1, 2 \quad (8.9)$$

where

$$\begin{bmatrix} u_1 & u_3 \\ u_2 & u_4 \end{bmatrix} = \begin{bmatrix} 1 & -\lambda \\ 0 & 1 \end{bmatrix} \mathbf{W} \mathbf{S}^{-1} . \quad (8.10)$$

We define $\mathbf{W}$ to be given by (8.6), and the $\mathbf{L}_{2p}, \mathbf{L}_j$ are defined by the long formulae of Lemma 6.3. We evaluate all these formulae (mod $cD\Gamma$). Since

$$\log cD\Gamma = O(\log \|E\|) ,$$

so we can use binary numbers with $O(\log \|E\|)$ digits throughout. The long formulae for $\mathbf{V}_k$ in Lemma 6.3 are evaluated successively. Evaluating each formula (6.20) (mod $cD\Gamma$) takes $O(M(\log \|E\|))$ operations. We next must check in (6.23) that given $\mathbf{B}_0, \mathbf{B}_{k+1}, \mathbf{V}_i$ and $\mathbf{V}_j$ (mod $cD\Gamma$) we can calculate $\mathbf{v}_{k+1}$ (mod $cD\Gamma$). It is straightforward to calculate $\mathbf{B}_0(\mathbf{V}_{k_i} \otimes \mathbf{V}_{k_2})$. We use the unimodularity condition of the matrix $\mathbf{B}_{k+1}$, that the greatest common divisor of its 2×2 submatrices $\Delta_{ij} = \begin{bmatrix} b_{1i} & b_{1j} \\ b_{2i} & b_{2j} \end{bmatrix}$ is 1. By an algorithm similar to step 1 of Lagarias [20, Theorem 5.4], repeatedly using the Euclidean algorithm with the $\det(\Delta_{ij})$ we can find a factorization

$$cD\Gamma = m_{12}m_{13}m_{14}m_{23}m_{24}m_{34} \quad (8.11)$$

with the m_{ij} pairwise relatively prime and with

$$(m_{ij}, \det(\Delta_{ij})) = 1. \quad (8.12)$$

for all i, j . This takes $O(M(\log \|E\|) \log \|E\|)$ operations. (Alternatively we can guess a set of m_{ij} and check that they have the required properties.) Then

$$(\Delta_{ij})^{-1} \equiv (\det(\Delta_{ij}))^{-1} \begin{bmatrix} b_{2j} & -b_{1j} \\ -b_{2i} & b_{1i} \end{bmatrix} \pmod{m_{ij}} \quad (8.13)$$

and $(\det \Delta_{ij})^{-1} \pmod{m_{ij}}$ is calculated in $O(M(\log \|E\|) \log \|E\|)$ operations using [20, Corollary 3.4]. Hence

$$\mathbf{V}_{k+1} \equiv (\Delta_{ij})^{-1} [\mathbf{B}_0(\mathbf{V}_{k_1} \otimes \mathbf{V}_{k_2})]_{ij} \pmod{m_{ij}}, \quad (8.14)$$

where $[\mathbf{M}]_{ij}$ denotes the submatrix obtained taking columns i and j , yields $\mathbf{V}_{k+1} \pmod{m_{ij}}$. Finally we use the Chinese remainder theorem on each entry of $\mathbf{V}_{k+1}$ separately to obtain $\mathbf{V}_{k+1} \pmod{cD\Gamma}$ in $O(M(\log \|E\|) \log \|E\|)$ operations, by [20, Prop. 3.6]. Thus we may at last obtain $\mathbf{L}_{2p}, \mathbf{L}_j \pmod{cD\Gamma}$ in $O(M \log \|E\| (\log \|E\|)^3)$ operations, by (6.26). Next we calculate $(\mathbf{L}_{2p})^2, (\mathbf{L}_{2p})^4$ etc. by successive squarings and reductions (mod $cD\Gamma$), and use the binary expansion of k to evaluate $W \pmod{cD\Gamma}$ using (8.6) in $O(M(\log \|E\|) (\log \|E\|)^2)$ operations, noting the bound (8.7). Finally (8.10) is evaluated (mod $cD\Gamma$) and then (8.9) verified in a further $O(M(\log \|E\|))$ operations. Thus (4.36) is verified in $O(M(\log \|E\|) (\log \|E\|)^3)$ elementary operations.

Finally we check that the sign conditions (4.37) of Lemma 4.2 (v) can be verified by evaluating the long formulae using floating-point arithmetic with floating-point integers maintaining $c_0(\log D)^3 = O((\log \|E\|)^3)$ binary digits in both the exponent and fraction parts, where c_0 is a sufficiently large absolute constant fixed once and for all as described below. Basic terminology and error estimates for floating-point computations are given in Appendix B. We say that a normalized floating-point number $\bar{x} = f2^e$ with $\frac{1}{2} \leq f < 1$ approximates x to accuracy s significant figures if

$$|\bar{x} - x| < 2^{e-s}. \quad (8.15)$$

(Here (e, f) is the representation of $\bar{x}$ used in the calculation.) We wish to show u_1 and u_2 are computed to accuracy at least 1 significant figure, which permits determination of their signs. Assuming for the moment this accuracy is proved, it is straightforward to estimate the total number of elementary operations involved in evaluating all the long formulae to be

$O(M(\log \|E\|)^3)(\log \|E\|)^2$ which is $O(M(\log \|E\|)(\log \|E\|)^4)$. Note here that in evaluating V_{k+1} by (6.23) that we merely pick an invertible Δ_{ij} , and use

$$\mathbf{V}_{k+1} = (\Delta_{ij})^{-1}[\mathbf{B}_0 \mathbf{V}_{k_1} \otimes \mathbf{V}_{k_2}]_{ij} \quad (8.16)$$

evaluated in floating-point, noting that

$$\log(\det(\Delta_{ij})) = O(\log \|E\|) \quad (8.17)$$

using (6.24).

It remains to estimate the loss of significant figures during the floating-point computations. The sources of loss of accuracy in floating-point computations are roundoff error, exponent overflow, exponent underflow in multiplication, and loss of accuracy in addition to two nearly equal numbers of opposite signs (e.g. this includes exponent underflow during addition as a special case).

By using $O(\log \|E\|)^3$ digits in the exponent part, we guarantee that exponent overflow never occurs. Indeed, only $O(\log \|E\|)$ binary digits are needed to represent the exponent part e of any entry of $\mathbf{W}$, since

$$e = O(\|E\|^{3/2}(\log \|E\|)^2) \quad (8.18)$$

by (4.41). It is easy to check that the bound (8.18) applies to any exponent of every element occurring in the long formulae, since the V_j 's are just various $\mathbf{L}_k$ with $1 \leq k \leq p$, to which the bounds (5.23), (5.25) apply. Now as long as the floating point calculations agree with the two entries of the long formulae to one significant figure, their exponents must agree within ± 1 and these calculated exponents will then satisfy (8.18) and exponent overflow cannot occur. This demonstrates that exponent overflow cannot occur unless all significant digits have first been lost due to the other three sources of error.

We next show exponent underflow during multiplications can never occur unless all significant digits have first been lost due to the remaining two sources of error. Indeed the entries of the matrices $\mathbf{V}_j$ in Lemma 6.3 are known a priori to be nonzero integers by Lemma 5.1(iii), except for $\mathbf{L}_i$ with $|i| < 2$ (and if these occur they may be placed at the beginning of the computation, which is done in fixed point as explained below). The entries of $\mathbf{W}$ are nonzero integers since $\mathbf{W} = \pm \mathbf{L}_j$ for some $|j| \geq 2$. We may suppose the entries of $\mathbf{U}$ are nonzero integers, for if some $u_j = 0$ then since $\mathbf{U} = \mathbf{Z}$ satisfies the hypotheses of Lemma 4.1 the inequalities (4.9), (4.10) would imply the u_j are small enough that they could be calculated directly in fixed point as certificates in $O(M(\log \|E\|))$ operations to verify (iv), (v) of Lemma 4.2. Since these entries are nonzero integers, the exponents of their floating-point approximations must be ≥ 0 , and exponent underflow during multiplication cannot occur by Lemma B-1 in Appendix B. (We note that some multiplications by zero may occur, but these are exact using (B-16), (B-17) of Appendix B.)

We must now bound the effects of roundoff error and that of addition of nearly equal quantities of opposite signs. We start with $p = c_0(\log \|E\|)^3$ significant digits of accuracy. We first consider the calculation of the $\mathbf{V}_k$ in Lemma 6.2. The entries of $\mathbf{S}_k, \mathbf{B}_0, \mathbf{B}_{k+1}$ are known to p significant digits by the bounds (6.21), (6.24). We will use Lemma B-2 to bound roundoff error, and Corollary B-4 to bound addition of nearly equal quantities. Evaluating $\mathbf{V}_{k+1}$ by the long formula (6.20) involves a loss of at most 5 significant digits by Lemma B-1, since each entry of $\mathbf{V}_{k+1}$ uses two floating-point multiplications and one addition, and the quantities

added always have the same sign by Lemma 5.1 (i), (ii). The crucial step lies in showing that evaluating $\mathbf{V}_{k+1}$ by the long formula (6.23) (actually by (8.16) above) involves a loss of at most $0(\log D)$ significant digits accuracy. Indeed $\mathbf{V}_{k_1} \otimes \mathbf{V}_{k_2}$ can be evaluated losing at most 3 significant digits accuracy by Lemma B-1, as only multiplications are involved. Now the bound (6.17) applies to show that

$$\|\mathbf{V}_k\| \geq \frac{2}{\|\mathbf{B}_k\|} (\|\mathbf{V}_{k_1}\| \|\mathbf{V}_{k_2}\|). \quad (8.19)$$

hence

$$\log \|\mathbf{V}_k\| \geq \log \|\mathbf{V}_{k_1}\| + \log \|\mathbf{V}_{k_2}\| - c_1 \log D - 1 \quad (8.20)$$

using (6.10). But all entries of $\mathbf{V}_k$ have about the same size by Lemma 5.1 (iii), hence the nearest floating-point approximations to each entry of $\mathbf{V}_k$ must have exponents e satisfying

$$e \geq \log \|\mathbf{V}_{k_1}\| + \log \|\mathbf{V}_{k_2}\| - (c_1 + 2) \log D - 3. \quad (8.21)$$

On the other hand, each entry of $\mathbf{V}_{k_1} \otimes \mathbf{V}_{k_2}$ has exponent

$$e \leq \log \|\mathbf{V}_{k_1}\| + \log \|\mathbf{V}_{k_2}\|. \quad (8.22)$$

We now evaluate the entries of (8.16) doing all multiplications first, followed by additions. The multiplications lose at most 6 significant digits each, and the resulting exponents satisfy

$$e \leq \log \|\mathbf{V}_{k_1}\| + \log \|\mathbf{V}_{k_2}\| + 2c_1(\log D) + 3 \quad (8.23)$$

using (8.22). Then the additions producing a given entry of $\mathbf{V}_{k+1}$ lose at most

$$(3c_1 + 2) \log D + 11$$

significant digits accuracy, using Corollary B-4, using (8.23) as an upper bound on e and (8.21) as a lower bound on $e - A$. Thus at most $(8c_1 + 2) \log D + 17$ significant digits are lost in evaluating $\mathbf{V}_{k+1}$ using the long formula (6.23), and thus at most $O((\log \|E\|)^3)$ significant digits are lost in evaluating $\mathbf{L}_{2p}$ and $\mathbf{L}_j$ using Lemma 6.2. Next, we note that the calculation of $(\mathbf{L}_{2p})^k$ in formula (8.6) involves a loss of $9(\log D)$ significant digits, because $O(\log D)$ matrix multiplications are involved in computing $(\mathbf{L}_{2p})^2, (\mathbf{L}_{2p})^4$ etc., and the bounds of Lemma B-2 apply because all numbers added have the same sign. Calculating $\mathbf{W}$ using the long formula (8.6) loses another 5 significant digits; again all quantities added have the same sign. Finally we evaluate

$$\mathbf{U} = \begin{bmatrix} 1 & -\lambda \\ 0 & 1 \end{bmatrix} \mathbf{W} \mathbf{S}^{-1}. \quad (8.24)$$

where $\lambda = \lceil \sqrt{D} \rceil$. Now

$$\mathbf{W} = \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix} \mathbf{U} \mathbf{S}$$

so

$$\|\mathbf{W}\| \leq 2\sqrt{D} \|\mathbf{U}\| \|\mathbf{S}\|$$

yields

$$\log \|\mathbf{U}\| \geq \log \|\mathbf{W}\| - c_2 \log \|\mathbf{E}\|. \quad (8.25)$$

for some absolute constant c_2 , using (4.40). The exponents e of the individual entries of $\mathbf{U}$ all satisfy

$$e \geq \log \|\mathbf{W}\| - c_3 \log \|\mathbf{E}\| \quad (8.26)$$

using the inequality (4.10) of Lemma 4.1 (which applies since $\mathbf{U} = \mathbf{Z}$). Now evaluate the right side of (8.24), doing all multiplications first, and then additions. The resulting multiplied quantities all have exponents

$$e \leq \log \|\mathbf{W}\| + (c_1 + 2) \log \|\mathbf{E}\| + 3 . \quad (8.27)$$

Then Corollary B-4 guarantees we can evaluate $\mathbf{U}$ with a loss of at most $(c_1 + c_3 + 2) \log \|E\| + 7$ significant digits accuracy. We have shown at most $O((\log \|E\|)^3)$ significant digits accuracy can be lost due to roundoff and adding of nearly equal quantities of opposite sign in evaluating $\mathbf{U}$. Choosing c_0 large enough once and for all, Theorem 1.1 is proved. $\square$

Proof of Theorem 1.2. All that need be checked is that the certificates of Theorem 1.1 can be “guessed” in polynomial time. The bounds (vi)–(ix) of Lemma 4.2, the bounds on k in Lemma 5.3 and on the $\mathbf{S}_{k+1}, \mathbf{B}_{k+1}$ in Lemma 6.3 demonstrate that this is the case. $\square$

Appendix A. Period lengths (mod m) of certain linear recurrences.

Let (t_1, u_1) be the least strictly positive solution to Pell's equation

$$X^2 - DY^2 = 1 \quad (\text{A.1})$$

and set

$$\epsilon = t_1 + u_1\sqrt{D} . \quad (\text{A.2})$$

In this appendix we show the sequences $\{t_k\}, \{u_k\}$ defined by

$$(\epsilon)^k = t_k + u_k\sqrt{D} \quad (\text{A.3})$$

are periodic (mod m) and we bound the length of the minimal period $P(m)$ for which

$$t_{k+p(m)} \equiv t_k \pmod{m} \quad (\text{A.4})$$

$$u_{k+p(m)} \equiv u_k \pmod{m} \quad (\text{A.5})$$

both hold.

The sequences $\{t_k\}, \{u_k\}$ both satisfy the second order linear recurrence.

$$w_k = t_1 w_{k-1} - w_{k-2} . \quad (\text{A.6})$$

Periodicity of solutions to this recurrence (mod m) is closely related to divisibility of u_k by m . Carmichael [9], [10] studied divisibility properties of a class of sequences which includes $\{t_k\}, \{u_k\}$ as special cases. Periodicity properties for general linear recurrences were considered by Engstrom [14], Ward [38] and other authors.

Lemma A-1. *The period $p(m)$ exists.*

Proof. Let $\bar{\epsilon} = t_1 - u_1\sqrt{D}$ so that

$$t_k = \frac{1}{2}(\epsilon^k + \epsilon^{-k}) \quad (\text{A.7})$$

$$u_k = \frac{1}{2\sqrt{D}}(\epsilon^k - \epsilon^{-k}) . \quad (\text{A.8})$$

Pell's equation asserts that

$$\epsilon\bar{\epsilon} = 1 . \quad (\text{A.9})$$

Thus $\epsilon, \bar{\epsilon}$ are units in the ring of integers O_D of $Q(\sqrt{D})$. For any ideal σ in θ_D let $S(\sigma)$ denote the smallest k such that

$$\epsilon^k \equiv \bar{\epsilon}^{-k} \equiv 1 \pmod{\sigma} \quad (\text{A.10})$$

over O_D , such $S(\sigma)$ existing since $\epsilon, \bar{\epsilon}$ are invertible (mod σ). It's easy to check that

$$t_{k+R} \equiv t_k \pmod{m}$$

$$u_{k+R} \equiv u_k \pmod{m}$$

where $R = S((2\sqrt{D}m))$. Hence

$$P(m) | S((2\sqrt{D}m)) \quad (\text{A.11})$$

exists. $\square$

Lemma A-2. *If $(m, n) = 1$ then*

$$p(mn) = l.c.m.\{p(m), p(n)\} . \quad (\text{A.12})$$

Proof. This follows from the definition (A-4) and the Chinese Remainder Theorem. $\square$

It thus suffices to calculate $P(p^a)$ for prime powers p^a .

Lemma A-3. *For all primes p and $a \geq 1$,*

$$P(p^{a+i}) | pP(p^a) . \quad (\text{A.13})$$

Proof. For $R = P(p^a)$ we have

$$\begin{aligned} t_R &= 1 + p^a \circ s_1 \\ u_R &= p^a s_2 \end{aligned} \quad (\text{A.14})$$

for some s_1, s_2 . Since

$$t_{pR} + u_{pR}\sqrt{D} = (t_R + u_R\sqrt{D})^p$$

we have

$$t_{pR} = \sum_{j=0}^{p-1/2} \binom{P}{2j} (t_R)^{p-2j} (u_R)^{2j} D^j \quad (\text{A.15})$$

$$u_{pR} = \sum_{j=0}^{p-1/2} \binom{P}{2j} (t_R)^j (u_R)^{p-2j} D^{\frac{p-1}{2}-j} \quad (\text{A.16})$$

Since $p | \binom{P}{j}$ for $1 \leq j \leq p-1$, these equations and (A.14) yield

$$t_{pR} \equiv (t_R)^p \equiv 1 \pmod{p^{a+1}}$$

$$u_{pR} \equiv 0 \pmod{p^{a+1}} .$$

and (A.13) follows. $\square$

In order to bound $P(p)$, let $(\frac{D}{p})$ denote the Legendre symbol and $Q(p)$ denote $Q((p))$.

Lemma A-4. *Let p be an odd prime.*

(i) *If $(\frac{D}{p}) = 1$, then*

$$P(p) | p-1 \quad (\text{A.17})$$

(ii) If $(\frac{D}{p}) = -1$, then

$$P(p)|2(p+1) . \quad (\text{A.18})$$

(iii) If $p|D$, then

$$P(p)|2p . \quad (\text{A.19})$$

(iv) $P(2) = 1$ or 2 .

Proof. Suppose $p \nmid 2D$ so $(\frac{D}{p}) = \pm 1$. Then examination of (A.7)–(A.10) shows that (A.11) can be sharpened to

$$P(p)|S(p) . \quad (\text{A.20})$$

(i) If $(\frac{D}{p}) = 1$, then (p) factors as $(p) = \gamma_1 \gamma_2$ the product of two distinct conjugate prime ideals in O_D . Then $O_D/\gamma_i \cong GF(p)$. Since $x^{p-1} = 1$ in $GF(p)$ when $x \neq 0$, we have

$$\epsilon^{p-1} \equiv \bar{\epsilon}^{p-1} \equiv 1 \pmod{\gamma_i} .$$

for $i = 1, 2$. Thus

$$\epsilon^{p-1} \equiv \bar{\epsilon}^{p-1} \equiv 1 \pmod{(p)}$$

so $S(p)|p-1$. Then (A.20) proves (A.17).

(ii) If $(\frac{D}{p}) = -1$, then (p) is inert, and $O_D/(p) \cong GF(p^2)$. Now $x^{p+1} \in GF(p)$ for all $x \in GF(p^2)$ hence

$$\epsilon p + 1 \equiv a \pmod{(p)} .$$

for some $a \in Z$. (Note $GF(p) \cong Z/pZ \subseteq O_D/(p)$.) Applying the conjugation automorphism, we have

$$\bar{\epsilon}^{p+1} \equiv a \pmod{(p)} .$$

But $\epsilon \bar{\epsilon} = 1$ hence

$$a^2 \equiv 1 \pmod{(p)} .$$

Hence

$$\epsilon^{2(p+1)} \equiv \bar{\epsilon}^{2(p+1)} \equiv 1 \pmod{(p)}$$

and $S(p)|2(p+1)$. Then (A.20) implies (A.18).

(iii) If $p|D$ then

$$t_2 = t_1^2 - Du_1^2 \equiv t_1^2 \equiv 1 \pmod{p}$$

since $t_1^2 = 1 + Du_1^2$. Then (A.15), (A.16) applied with $R = 2$ show

$$t_{2p} \equiv (t_2)^p \equiv 1 \pmod{p}$$

$$u_{2p} \equiv 0 \pmod{p} .$$

Hence $P(p)|2p$.

(iv) If $2|D$ then $t_1^2 - Du_1^2 = 1$ shows $t_1^2 \equiv 1 \pmod{4}$ and $2|u$, or $4|D$. In either case $t_2 \equiv 1 \pmod{4}$, $u_2 \equiv 0 \pmod{2}$ and $P(2)|2$. If $2 \nmid D$ then $(t_1, u_1) \equiv (1, 0)$ or $(0, 1) \pmod{2}$. In the first case $P(2) = 1$, in the second case, the recurrence (A.15) shows $P(2) = 2$. $\square$

Lemma A-5. *For any m ,*

$$P(m) \leq 2m(\log m + 1)$$

Proof. Lemmas A-2 through A-4 imply that if $m = \prod_j (p_j)^{a_j}$ then $P(M)|R(M)$ where

$$R(m) = 2\prod_j p_j^{a_j-1} \left(p_j - \left(\frac{D}{p} \right) \right) .$$

Now

$$R(m) \leq 2m\prod_j \left(1 + \frac{1}{p_j} \right) .$$

But

$$\pi_j \left(1 + \frac{1}{p_j} \right) \leq \sum_{j=1}^m \frac{1}{j} < \log m + 1 \tag{A.21}$$

and the lemma follows. $\square$

Remark. By more detailed argument it can be shown that

$$P(m) = O(m \log \log m) .$$

Appendix B. Floating-Point Computations.

This appendix gives upper bounds on the magnitude of errors accumulated in floating-point computations. We use the conventions and notation of Knuth [14, Sect. 4.2], to which we refer for greatest detail.

We use *normalized floating-point numbers* with *base 2*, *excess 0*, with *p digits*. Such a number will be denoted (e, f) where

$$(e, f) = f2^e . \quad (\text{B.1})$$

Here e is an integer satisfying

$$|e| < N \quad (\text{B.2})$$

and f is a signed fraction such that $2^p f$ is an integer and satisfying the *normalization condition*

$$\frac{1}{2} \leq |f| < 1 . \quad (\text{B.3})$$

provided $f \neq 0$. By convention 0 is (0,0).

We introduce a notation to distinguish general real numbers from floating-point numbers, which are just real numbers satisfying (B.1)–(B.3). To this end we always denote floating-point numbers with a bar, i.e., $\bar{x}$ is a floating-point number (to be thought of as an approximation to the real number x).

To define the floating-point operations of addition, subtraction, multiplication and division, we use the function “Round to p significant figures” defined by

$$\text{Round } (x, p) = \begin{cases} 2^{e-p} \lfloor 2^{p-e} x + \frac{1}{2} \rfloor, & 2^{e-1} \leq x \leq 2^e \\ 0, & x = 0 \\ 2^{e-p} \lceil 2^{p-e} x - \frac{1}{2} \rceil, & 2^{e-1} \leq -x < 2^e . \end{cases} \quad (\text{B.4})$$

We define *floating-point addition* $\oplus$ by

$$\bar{x} \oplus \bar{y} = \begin{cases} 0, & |\bar{x} + \bar{y}| < 2^{-N} \\ \text{Round } (x + y, p), & 2^{-E} \leq |\bar{x} + \bar{y}| < 2^N . \end{cases} \quad (\text{B.5})$$

Exponent overflow occurs if $|\bar{x}\bar{y}| \geq 2^N$ and $\bar{x} \oplus \bar{y}$ is left undefined. We define floating-point subtraction of $\bar{x}$ as floating-point addition of $-\bar{x}$. We define *floating-point multiplication* $\otimes$ by

$$\bar{x} \otimes \bar{y} = \begin{cases} 0, & |\overline{xy}| < 2^{-N} \\ \text{Round } (\overline{xy}, p), & \text{if } 2^{-E} \leq |\overline{xy}| < 2^N \end{cases} \quad (\text{B.6})$$

Exponent overflow occurs if $|\overline{xy}| \geq 2^N$ and $\bar{x} \otimes \bar{y}$ is left undefined. Floating-point division ϕ is defined similarly to multiplication, but we will not need it. Note that these operations are well-defined even when *exponent underflow* occurs.

Let $\bar{x}$ be a floating point number approximating a nonzero real number x . Let

$$2^e \leq x < 2^{e+1} . \quad (\text{B.7})$$

We say $\bar{x}$ *approximates* x to s significant digits if

$$|\bar{x} - x| < 2^{e-s-1} . \quad (\text{B.8})$$

There are four sources of loss of significant digits in floating-point operations.

1. roundoff error,
2. exponent overflow,
3. exponent underflow in multiplication,
4. addition of two nearly equal quantities of opposite signs (includes exponent underflow).

We deal with these sources separately.

Exponent overflow, and exponent underflow in multiplication are the easiest to handle, by giving sufficient conditions that they do not occur. By convention multiplication by zero does not count as exponent underflow.

Lemma B-1. *Let $\bar{x} = (e_1, f_1)\bar{y} = (e_2, f_2)$ be two floating-point numbers. If*

$$-N + 2 \leq e_1 + e_2 \leq N - 1 \quad (\text{B.9})$$

then $\bar{x} \otimes \bar{y}$ does not involve exponent overflow or underflow. If

$$\text{MAX}(e_1, e_2) \leq N - 2 \quad (\text{B.10})$$

then $\bar{x} \oplus \bar{y}$ does not involve exponent overflow.

Proof. Immediate. $\square$

In the remainder of this Appendix we assume that exponent overflow does not occur. That is, this is an extra hypothesis made in all lemmas following.

In order to analyze roundoff error, we note that when

$$2^{e-1} \leq |x| < 2^e \quad (\text{B.11})$$

we have the bound

$$|\text{Round}(x, p) - x| < 2^{e-p-1} . \quad (\text{B.12})$$

Lemma B-2. *Let $\bar{x} - \bar{y}$ be two floating-point numbers, both having s significant digits.*

- (i) *If $\bar{x}, \bar{y}$ have the same sign, then at most 2 significant digits are lost in computing $\bar{x} \oplus \bar{y}$.*
- (ii) *If exponent underflow does not occur, at most 3 significant digits are lost in computing $\bar{x} \otimes \bar{y}$.*

Proof. (i) Since $\bar{x}, \bar{y}$ have the same sign, underflow cannot occur. Then

$$\bar{x} \oplus \bar{y} = \text{Round} \quad (\bar{x} + \bar{y}, p) . \quad (\text{B.13})$$

Let $\bar{x}, \bar{y}$ have exponents e_1, e_2 . Then the exponent e_3 of $\bar{x} \oplus \bar{y}$ is at least $\text{MAX}(e_1, e_2)$. But

$$\begin{aligned} |\bar{x} - x| &< 2^{e_1-s-1} \\ |\bar{y} - y| &< 2^{e_2-s-1} . \end{aligned} \quad (\text{B.14})$$

Note $s \leq p$. Then

$$\begin{aligned} |\bar{x} \oplus \bar{y} - (x + y)| &\leq |\bar{x} \oplus \bar{y} - (\bar{x} + \bar{y})| + |\bar{x} - x| + |\bar{y} - y| \\ &\leq 2^{e_3-p-1} + 2^{e_2-s-1} + 2^{e_1-s-1} \leq 2^{e_3-s+1} \end{aligned} \quad (\text{B.15})$$

using (B.12), (B.13).

(ii) Since underflow cannot occur, we have

$$\bar{x} \otimes \bar{y} = \text{Round} \quad (\overline{xy}, p) . \quad (\text{B.16})$$

If e_4 is the exponent of $\bar{x} \otimes \bar{y}$, then

$$e_4 \geq e_1 + e_2 - 1 .$$

Now

$$|\overline{xy} - xy| \leq |\bar{x} - x| |\bar{y}| + |\bar{y} - y| |x| \leq 2^{e_1+e_2-s}$$

using (B.14). Hence

$$\begin{aligned} |\bar{x} \otimes \bar{y} - xy| &\leq |\bar{x} \otimes \bar{y} - \overline{xy}| + |\overline{xy} - xy| \\ &\leq 2^{e_3-p-1} + 2^{e_1+e_2-s} \leq 5 \cdot 2^{e_3-s-1} , \end{aligned}$$

using (B.12), (B.16). $\square$

We remark that Lemma B-2 (i) also holds when $\bar{y} = 0$ and

$$|\bar{y} - y| < 2^{e_1-s-1} , \quad (\text{B.17})$$

and that

$$\bar{x} \otimes \bar{y} = xy = 0 \quad (\text{B.18})$$

where $y = \bar{y} = 0$.

We next consider the bounds for addition.

Lemma B-3. *Let $\bar{x}_1, \dots, \bar{x}_j$ be floating-point numbers such that all $\bar{x}_i$ have exponents $\leq e$. Suppose*

$$|\bar{x}_i - x_i| < 2^{e-s-1} \quad (\text{B.19})$$

for all i , and suppose $e - s \geq -N$. Let

$$v_j = x_1 + \dots + x_j , \quad (\text{B.20})$$

and define $\bar{v}_1 = \bar{x}_1$ and

$$\bar{v}_{i+1} = \bar{v}_i \oplus \bar{x}_{i+1} \quad (\text{B.21})$$

for $2 \leq i \leq j-1$. Let $j \leq 2^k$ and $k \leq p$. Then

$$|\bar{v}_j - v_j| < 2^{3+2k+3-s} . \quad (\text{B.22})$$

Proof. We have

$$|\bar{x}_i| \leq 2^e - 2^{e-p} ,$$

from which it is easy to establish

$$\bar{v}_i \leq i(2^e - 2^{e-p})(1 + i2^{-p}) < 2^{e+k+2} . \quad (\text{B.23})$$

(The term $i2^{-p}$ is a roundoff bound.) Now we have

$$|\bar{v}_i - v_i| \leq |\bar{v}_i - (\bar{v}_{i-1} + \bar{x}_i)| + |\bar{x}_i - x_i| + |\bar{v}_{i-1} - v_{i-1}| . \quad (\text{B.24})$$

If we let e_i be the exponent of $\bar{v}_i$ then (B.22) gives

$$e_i \leq e + k + 2 . \quad (\text{B.25})$$

But

$$|\bar{v}_i - (\bar{v}_{i-1} + \bar{x}_i)| \leq \text{MAX}(2^{e_i-p-1}, 2^{-N}) , \quad (\text{B.26})$$

the bound 2^{-N} occurring in the case of underflow. Then apply (B-18) and (B-25) to (B-22) and sum over i to obtain

$$|\bar{v}_j - v_j| < \sum_{i=1}^j [2^{e_j-p-1} + 2^{-N} + 2^{e-s-1}] . \quad (\text{B.27})$$

using (B-24) gives

$$\begin{aligned} |\bar{v}_j - v_j| &< 2^{e+2k+1-p} + 2^{N+k} + 2^{e+k-s-1} \\ &< 2^{e+wk+3-s} , \end{aligned}$$

the desired bound. $\square$

Lemma B-3 allows one to show that if one knows “a priori” that a sum $\sum_i x_i$ is not too small with respect to its largest term, then the loss of significant digits in calculating a floating-point approximation to this sum cannot be large.

Corollary B-4. *Let $\bar{x}_1, \dots, \bar{x}_j$ be floating point numbers approximating $x_1, \dots, x_j$ to s significant digits, with the largest $|\bar{x}_j|$ having exponent $e \geq -N + s$. Let*

$$s_j = x_1 + \dots + x_j ,$$

$$\bar{s}_j = \bar{s}_{j-1} \oplus \bar{x}_j, \bar{s}_1 = \bar{x}_1 ,$$

Suppose $j \leq 4$ and that

$$|s_j| \geq 2^{e-A} .$$

Then $\bar{s}_j$ approximates s_j to at least $s - A - 8$ significant digits. $\square$

References

- [1] L. Adleman, Number Theoretic Aspects of Computational Complexity, Thesis, Univ. of California, Berkeley, 1976.
- [2] L. Adleman and K. Manders, Diophantine Complexity, Proc. 17th IEEE Annual Symp. on Foundations of Computer Science (1976), 81–88.
- [3] L. Adleman and K. Manders, Reducibility, Randomness and Intractability, Proc. 9th Annual ACM Symposium on Theory of Computing (1977), 151–163.
- [4] L. Adleman and K. Manders, Intractability Proofs and the Computational Complexity of Binary Quadratics, U. C. Berkeley, College of Engineering Technical Report No. UCB/ERL M78/30 (1978).
- [5] L. Adleman and K. Manders, Reductions that Lie, Proc. 20th IEEE Symp. on Foundations of Computer Science (1979), 397–410.
- [6] M. Agrawal, N. Kayal and N. Saxena, PRIMES is in P, *Annals of Math.* **160** (2004), 781–793.
- [7] J. Buchmann and H. C. Williams, On the existence of a short proof for the value of the class number and the regulator of a real quadratic field, in: *Number Theory and Applications (Banff, AB 1988)*, 327–345, NATO ASI Series C Math. Phys. Sci, No. 265, Kluwer Academic Publ., Dordrecht 1989.
- [8] D. Buell, *Binary Quadratic Forms*. Classical theory and modern computations. Springer-Verlag, New York 1989.
- [9] R. D. Carmichael, On the numerical factors of the arithmetic forms $\alpha^n \pm \beta^n$, *Annals of Math.* **15** (1913), 30–70.
- [10] R. D. Carmichael, A Simple Principle of Unification in the Elementary Theory of Numbers, *Amer. Math. Monthly* **36** (1929), 132–143.
- [11] H. Cohen, *A Course in Computational Algebraic Number Theory*, Graduate Texts in Mathematics 138, Springer-Verlag: New York 1993.
- [12] M. Davis, Hilbert’s Tenth Problem is Unsolvable, *Amer. Math. Monthly* **80** (1973), 233–269.
- [13] M. Davis, H. Putnam and J. Robinson, The Decision Problem for Exponential Diophantine Equations, *Annals of Math.* **74** (1961), 425–436.
- [14] H. T. Engstrom, On Sequences Defined by Linear Recurrence Relations, *Trans. Amer. Math. Soc.* **33** (1931), 210–218.
- [15] C. F. Gauss, *Disquisitiones Arithmeticae 1801*. (English translation: Yale U. Press, New Haven, 1966.)

- [16] E. M. Gurari and O. H. Ibarra, An *NP*-Complete Number-Theoretic Problem, *J. ACM* *26* (1979), 567–581.
- [17] L. K. Hua, On the least solution to Pell’s equation, *Bull. Amer. Math. Soc.* *48* (1942), 731–735.
- [18] D. E. Knuth, *The Art of Computer Programming, Vol. 2, Seminumerical Algorithms*, Addison-Wesley Publ. Co., Reading, Mass. 1969.
- [19] J. C. Lagarias, Succinct Certificates for the Solvability of Binary quadratic Diophantine Equations (Extended Abstract), *Proc. 20th IEEE Symp. on Foundations of Computer Science* (1979), 47–54.
- [20] J. C. Lagarias, Worst-case complexity bounds for algorithms in the theory of integral quadratic forms, *J. of Algorithms* *1* (1980), 42–86.
- [21] J. C. Lagarias, On the computational complexity of determining the solvability or unsolvability of the equation $x^2 - Dy^2 = -1$, *Trans. Amer. Math. Soc.* *260* (1980), 485–508.
- [22] J. C. Lagarias, Succinct Certificates for the Solvability of Binary quadratic Diophantine Equations, Bell Labs Technical Memorandum 81-11216-54, Sept. 28, 1981.
- [23] H. W. Lenstra, Jr, On the calculation of regulators and class numbers of quadratic fields, in: *Number theory days, 1980 (Exeter 1980)*, pp. 123–150, London Math. Soc. Lecture Notes Series 56, Cambridge University Press, Cambridge 1982.
- [24] K. Manders and L. Adleman, *NP*-complete decision problems for binary quadratics, *J. Comp. Sys. Sci.* *16* (1978), 168–184.
- [25] G. B. Mathews, *Theory of Numbers*, 2nd Ed., New York, 1961 (Reprint).
- [26] Y. Matijasevic, Enumerable Sets are Diophantine, *Dokl. Akad. Nauk SSSR* *191* (1970), 279–282.
- [27] G. Miller, Riemann’s Hypothesis and Tests for Primality, *J. Computer and Systems Science* *13* (1976), 300–317.
- [28] W. Narkiewicz, *Elementary and Analytic Theory of Algebraic Numbers*, Polish Scientific Publishers, Warsaw, 1974.
- [29] V. Pratt, Every Prime has a Succinct Certificate, *SIAM J. Computing* *4* (1975), 214–220.
- [30] L. Redei, Die 2-Ringklassengruppe des Quadratischen Zahlkörperws und die theorie der Pellschen Gleichung, *Acta. Math. Acad. Sci. Hung.* *4* (1953), 31–87.
- [31] R. L. Rivest, A. Shamir and L. Adleman, A method for obtaining digital signatures and public key cryptosystems, *Comm. ACM* *21* (1978), 120–126.
- [32] D. Shanks, Class number, a theory of factorization and genera, in: 1969 Number Theory Institute, *Proc. Symp. Pure Math. XX* (1971), 415–440.

- [33] D. Shanks, The Infrastructure of a Real Quadratic Field and Its Applications, Proc. 1972 Number Theory Conf., U. of Colorado, Boulder, Colorado (1972), 217–224.
- [34] S. Smale, Mathematical problems for the next century, in: *Mathematics: frontiers and perspectives*, pp. 271–294, Amer. Math. Soc., Providence RI 2000.
- [35] H. J. S. Smith, *Report on the Theory of Numbers*, Chelsea Publ. Co., New York (Reprint).
- [36] C. Thiel, Under the assumption of the generalized Riemann hypothesis, verifying the class number belongs to $\text{NP} \cap \text{co-NP}$, in: *Algorithmic number theory (Ithaca, NY 1994)*, pp. 234–247, Lecture Notes in Computer Science 877, Springer-Verlag, Berlin 1994.
- [37] B. A. Venkov, *Elementary Number Theory*, Wolters-Noordhoff Publ. Co., Groningen, The Netherlands, 1970.
- [38] M. Ward, The arithmetical theory of linear recurring series, Trans. Amer. Math. Soc. 35 (1933) 600–628.